Appendix F. Flora and Fauna Assessment

ADVERTISED PLAN

This copied document to be made available for the sole purpose of enabling its consideration and review as part of a planning process under the Planning and Environment Act 1987. The document must not be used for any purpose which may breach any convright

Jacobs

Prospect Hill EfW Project

Qualified Desktop Flora and Fauna Assessment

1 | 2 2 June 2020

Prospect Hill International Pty Ltd

Α

Document history and status

| Revision | Date | Description | Author | Checked | Reviewed | Approved |
|----------|------------|------------------------|------------|---------|------------|------------|
| 1 | 06/05/2020 | Edits to DRAFT | D Endersby | | A Stephens | D Endersby |
| 2 | 02/06/2020 | Site Assessment Update | D Endersby | | K Munro | D Endersby |
| | | | | | | |
| | | | | | | |
| | | | _ | | | |

Distribution of copies

| Revision | lssue approved | Date issued | Issued to | Comments |
|----------|-------------------|----------------|-----------|----------|
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |

Prospect Hill EfW Project

| Project No: | IS305100 |
|------------------|--|
| Document Title: | Qualified Desktop Flora and Fauna Assessment |
| Document No.: | 1 |
| Revision: | 2 |
| Date: | 2 June 2020 |
| Client Name: | Prospect Hill International Pty Ltd |
| Client No: | A |
| Project Manager: | Kate Munro |
| Author: | David Endersby |
| File Name: | R02_Prospect Hill EfW Plant_FF.docx |

Jacobs Australia Pty Limited

Floor 11, 452 Flinders Street Melbourne VIC 3000 PO Box 312, Flinders Lane Melbourne VIC 8009 Australia T +61 3 8668 3000 F +61 3 8668 3001 www.jacobs.com

© Copyright 2019 Jacobs Australia Pty Limited. The concepts and information contained in this document are the property of Jacobs. Use or copying of this document in whole or in part without the written permission of Jacobs constitutes an infringement of copyright.

Limitation: This document has been prepared on behalf of, and for the exclusive use of Jacobs' client, and is subject to, and issued in accordance with, the provisions of the contract between Jacobs and the client. Jacobs accepts no liability or responsibility whatsoever for, or in respect of, any use of, or reliance upon, this document by any third party.



Contents

| Execut | tive Summaryiii |
|--------|---|
| 1. | Introduction1 |
| 1.1 | Location1 |
| 2. | Method2 |
| 2.1 | Desktop assessment |
| 2.1.1 | Site assessment |
| 2.1.2 | Relevant legislation and policies |
| 3. | Desktop Result |
| 3.1 | Native Vegetation |
| 3.1.1 | Ecological Vegetation Class (EVC) |
| 3.1.2 | Listed Ecological Communities (FFG Act)7 |
| 3.1.3 | Threatened Ecological Communities (EPBC)8 |
| 3.1.4 | Native Vegetation Preprint Plan9 |
| 3.1.5 | Habitat9 |
| 3.1.6 | Site Assessment9 |
| 3.2 | Threatened Species9 |
| 3.2.1 | Threatened Species (PMST) |
| 4. | Assessment |
| 5. | Conclusion and recommendations |
| 6. | References |

Appendix A. Protected Matters Search Tool Report

Appendix B. PMST Threatened Species Assessment

Executive Summary

This qualified desktop flora and fauna assessment considers potential statutory flora and fauna issues associated with the site of the proposed Prospect Hill Energy from Waste (EfW) Project at164-200 McManus Road, Lara.

A range of flora and fauna data sets were reviewed, including the relevant Precinct Structure Plan (PSP) for the Geelong Ring Road Employment Precinct (GREP).

Desktop data reviewed from state and commonwealth agencies suggested a moderate likelihood of threatened species remaining in and around the site, however, a review of the relevant Native Vegetation Precinct Plan (NVPP) where it was determined that no native vegetation persisted on site, limits the likelihood of occurrence associated with most threatened species relevant to the area. A brief site assessment was undertaken to confirm the results of the desktop assessment against site attributes, particularly any habitat remaining, given the level of modification and disturbance the site has been subject to in recent times. This assessment found there to be no habitat remaining likely to be utilised by threatened species, and that, as reported in the PSP Native Vegetation Assessment, no native vegetation remains on site.

It is noted the site has been largely cleared since approvals were granted under the PSP process by City of Greater Geelong, resulting in the majority of habitat being removed from the site. Although a threatened species assessment was not conducted as part of the NVPP, there is some potential that a small number of threatened species may persist in the areas not subject to disturbance, particularly around the periphery of the site, although these areas are likely subject to various degradation issues.

Important note about your report

Information from the desktop assessment is only as reliable as the data available and in the case of the VBA the number of surveys previously undertaken (i.e. an area where many surveys have been taken in the past, will, most likely, have a more extensive list of species than areas where very little survey work has been undertaken). The accuracy of past surveys is also variable and point locations can be out by up to 1 km.

In addition to the number of previous surveys undertaken, there are other reasons why species, including threatened species, may not have previously been recorded. For example, at the time of historical site visits some plant species may not have been visible above the ground or flowering and therefore not identified as being present within the area surveyed. Also, the data collected is likely to consist of opportunistic observations only, and, therefore, listed fauna species moving in and out of the area may not have been observed or recorded. Similarly, many fauna species are cryptic, nocturnal and well-hidden such that their presence can only be detected through detailed targeted assessment methods. Hence on species that can be readily identified at that time, heard or have distinctive signs, such as tracks, scats, diggings are those most likely to be recorded.

The EPBC Act is undergoing a review that commenced in October 2019. Any changes to the applicable legislation and agreements may affect the outcomes of this report.

The FFG Act Amendment Bill 2019 has passed through Victorian Parliament with amendments taking effect on 1 June 2020. This report has been prepared based on the current requirements of the Act and these may change prior to the construction of the project.

Spatial data layers assessed were the most current available at the time of the assessment. Any changes to these layers may require the outcomes of this document to be updated.

1. Introduction

This qualified desktop flora and fauna assessment reviews the risks relating to statutory flora and fauna issues (including native vegetation) associated with the development of the Prospect Hill Energy from Waste Project (the Project).

This assessment reviews a number of existing databases of modelled and previously recorded species and communities from the wider landscape and considers the potential of such features to occur within the Project footprint, given site attributes and habitat apparent from the data reviewed.

This report provides a preliminary assessment to collate ecological values potentially occurring in the vicinity of the proposed Project.

1.1 Location

The Project site occurs within the City of Greater Geelong Local Government Area and the Corangamite Catchment Management Authority's (CMA) management area. The site occurs on the Victorian Volcanic Plain bioregion.

The assessment area is a small property approximately 16 ha in size at 164-200 McManus Road, Lara (Figure 1.1). Project location This site is north of the Geelong Ring Road and approximately 55 km south west of the Melbourne CBD.

The site has been significantly disturbed and modified through previous land use. The current state of the site appears to reflect recent storage of bulk earth materials, with all top soil and natural features apparently removed from the site as part of historical land management. The periphery of the site appears to be bunded, with a high density of weed species. A large proportion of the site appears to be cleared of all vegetation, with areas retaining vegetation appearing to be largely composed of grassy weeds.




Figure 1.1: Project location

2. Method

This report was compiled based on a desktop assessment and with a brief field qualification, as described in the following sections.

2.1 Desktop assessment

The desktop assessment involved a review of the following databases administered by the Victorian Department of Environment, Land, Water and Planning (DELWP) and Commonwealth Department of Agriculture, Water and the Environment (DAWE):

- Native Vegetation Mapping (DELWP, 2020a) This database comprises large-scale modelling and classification of native vegetation across Victoria. Information in the database includes modelled presence of native vegetation, modelled extent of Ecological Vegetation Classes (EVC) as of 2005, Strategic Biodiversity Score (indicative of the contribution of the area to Victoria's biodiversity values) and modelled condition of vegetation present.
- Victorian Biodiversity Atlas (VBA) (DELWP, 2020b) This database comprises historical records of flora and fauna species from across Victoria. Records are added opportunistically, as flora and fauna surveys are conducted within Victoria for a variety of purposes. The mapping of flora and fauna distribution and determination of species' habitat preferences is an ongoing process. Records from a 5 km buffer of the project area have been assessed for this report.
- Protected Matters Search Tool (PMST) (DAWE, 2020) The PMST highlights any Matters of National Environmental Significance (MNES) relevant to the *Environment Protection and Biodiversity Conservation* (*EPBC*) Act 1999 that are likely to occur within an area. Records from a 5 km buffer of the project area have been assessed for this report.
- Relevant Aerial Imagery Aerial imagery allows for high level habitat conditions to be assessed.
- Background documentation associated with the Precinct Structure Plan process and accompanying Native Vegetation Precinct Plan (Ecology Partners, 2011).

An assessment of the likelihood of threatened species and ecological communities occurring within and adjacent to the project area was then undertaken for species identified as potentially occurring during the database and literature searches. This assessment was based on the species' / communities' preferred habitat in comparison to the habitat available at the site, and the frequency, timing and location of previous recordings. This assessment focussed on species and ecological communities currently listed on Victoria's Advisory Lists of Threatened Species (VicAdv) (DEPI, 2013), or those listed under the *Flora and Fauna Guarantee Act 1988* (FFG Act) or EPBC Act. The criteria used for assessing likelihood of occurrence are described in Table 2.1 and Table 2.2.

A summary of the likelihood of occurrence assessment is provided in Section Table 2.1

| Table 2.1 : Criteria for determining the likelihood of threatened species being present within the project assessment |
|---|
| area. |

| Likelihood | Criteria |
|------------|--|
| High | Recent records of species from DELWP databases Review of aerial photography indicates potential habitat within the project area |
| | Review of habitat and distribution literature indicates the site is appropriate for this species. |
| Moderate | Historic records of species from DELWP databases Review of habitat distribution literature indicates the project area is appropriate for this species Review of aerial photography indicates limited habitat within the project area. |
| Low | Species has not been previously recorded within DELWP database Review of aerial photography indicates that no available habitat within the project area Review of literature regarding habitat and distribution indicates the project area is unlikely to be utilised by this species. |

Table 2.2 : Criteria for determining the likelihood of threatened ecological communities being present within the project assessment area

| Likelihood | Criteria |
|------------|---|
| High | Mapping by DELWP indicates the EVCs likely to be present at the site are of a similar composition to the threatened ecological community Review of aerial photography indicates that native vegetation is likely to be present at the site Review of literature and general knowledge of vegetation in the area indicates the site is appropriate for this threatened ecological community. |
| Moderate | Mapping by DELWP indicates the EVCs likely to be present at the site are of a similar composition to the threatened ecological community It is difficult to determine from aerial photography whether the community is present, such as grassland communities Review of literature and general knowledge of vegetation in the area indicates the project area is suitable for this threatened ecological community. |
| Low | Mapping by DELWP indicates the EVCs likely to be present at the site are not of similar composition to the threatened ecological community Review of aerial photography indicates that no native vegetation is likely present Review of literature and general knowledge of vegetation in the area indicates that the threatened ecological community is unlikely to be present at the project area. |

2.1.1 Site assessment

A brief site assessment was undertaken on the 29th of May to consider the desktop results derived from the state and federal databases detailed above, considered against the specific habitat features currently evident at the site. It is acknowledged site modification and disturbance that has occurred in recent times is not reflected in some of the data results provided, particularly site condition and native vegetation attributes presented in DELWP data.

2.1.2 Relevant legislation and policies

A brief summary of the legislation and policies referred to throughout the document is provided below in Table 2.3.

| Table | 2.3: | Relevant | legislation |
|-------|------|--------------------|-------------|
| 10010 | 2.0. | i i ci ci v ai i c | registation |

| Policy/legislation Description | | Project action | | | | | |
|--|--|--|--|--|--|--|--|
| Commonwealth | | | | | | | |
| Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act) | The EPBC Act has significant implications for natural resource and environmental management in Australia. This Act provides for the listing of threatened species, threatened ecological communities and key threatening processes. It also relates to actions likely to have a significant impact on Matters of National Environmental Significance (MNES), including World Heritage Sites National Heritage Places Ramsar Wetlands Nationally threatened species and ecological communities Migratory species Commonwealth marine areas Nuclear actions the Great Barrier Reef Marine Park | Determine whether MNES are likely to be 'significantly' impacted by the proposed works. Recommend further assessment as required, such as targeted surveys. Where MNES may be impacted recommend mitigation measures to avoid and reduce impact. If impact cannot be avoided the project will need to be referred to the Commonwealth Department of Agriculture, Water and the Environment (DAWE). | | | | | |

| Policy/legislation | Description | Project action |
|---|---|---|
| | a water resource, in relation to coal seam gas development or large coal mining development | |
| State | | |
| Environment Effects Act 1978 (EE Act) | The EE Act provides for the assessment of actions that are capable of having a significant environmental effect. Actions that may have a significant environmental effect should be referred to the Victorian Minister for Planning, who decides if an Environmental Effects Statement (EES) is required. An EES may be required where: 10 ha or more native vegetation is being cleared There is a likelihood of regionally or state significant adverse environmental effects There is a need for an integrated assessment of social and economic effects of a project or relevant alternatives Normal statutory processes would not provide a sufficiently comprehensive, integrated and transparent assessment This Act also allows an applicant to write to the Secretary of the Victorian Department of Environment Land Water and Planning (DELWP) to confirm no EES is required. | Determine whether the extent of removal of native vegetation and habitat for threatened species of state significance will trigger the need for a referral under the EE Act. Recommend further assessment where required, such as targeted surveys. If a trigger for EES is met, recommend mitigation measures to avoid and reduce impact. If impact cannot be avoided an EES referral will need to be submitted. |
| Flora and Fauna Guarantee Act 1988 (FFG Act) | The FFG Act provides a framework for biodiversity conservation in Victoria. Threatened species and communities of flora and fauna, as well as threatening processes, are Listed under this Act. A number of non-threatened flora species are also listed as protected under this Act. A Permit to Take is required to remove these species from public land. | Determine if Listed flora or fauna species are likely to be affected or threatening processes resultant of the proposed works. Recommend further assessment where required, such as target surveys. Where Listed flora and fauna species are identified or threatening processes likely, recommend mitigation measures to avoid and reduce impact. If Listed flora and fauna species are to be removed, a Permit to Take may need to be obtained. |
| DELWP Victorian Advisory Lists (VicAdv) | The DELWP Victorian Advisory Lists (VicAdv) are not a statutory list of threatened species, but rather list species for which conservation management is recommended by DELWP. The VicAdv lists are comprised of: the Advisory List of Rare or Threatened Plants in Victoria – 2014 (DEPI 2014) the Advisory List of Threatened Vertebrate Fauna in Victoria – 2013 (DEPI 2013), and the Advisory List of Threatened Invertebrate Fauna in Victoria – 2009 (DSE 2009). The presence, or likely presence, of a species listed on the VicAdv lists is used to determine whether species specific habitat is required to be offset. | Determine if species present are listed on the VicAdv lists and likely to be affected by the proposed works within the project area. Recommend further assessment where required, such as targeted surveys. Where listed flora and fauna species are identified, recommend mitigation measures to avoid and reduce impacts. If listed flora and fauna species are to be impacted an offset will be prescribed for the project area that incorporates habitat for the affected species. |
| Planning and Environment Act 1987 (P&E Act) | Applications to remove, destroy, or lop native vegetation in Victoria usually invoke Clause 52.17 of the Victorian Planning Provisions which is given authority through this | Determine if any planning overlays occur which require a permit for vegetation removal. |

Jacobs

| Policy/legislation | Description | Project action |
|---|---|---|
| | Act. Certain overlays within the planning scheme (e.g. Environmental Significance Overlays) may modify the permit requirements for the removal of native vegetation. A range of exemptions may apply. | Determine whether native vegetation is present and will be impacted by the project, requiring permitting, and consider whether exemptions may apply. |
| | Where a Precinct Structure Plan has been endorsed for an area, vegetation impacts are governed under Clause 52.16 of the P&E Act where the relevant council endorses | Review relevant PSP (Geelong Ring Road Employment Precinct (GREP) for any specified offset requirement. |
| Guidelines for the removal, destruction or lopping of native vegetation (the Guidelines) | development and offsets are managed through a relevant Vegetation Precinct Plan (NVPP). Permits to remove native vegetation are still required, however offsets are managed through the VMP process. | If native vegetation is to be impacted, a permit will be required from the approval authority and the appropriate offset requirements identified and obtained. |
| Catchment and Land Protection Act 1994 (CaLP Act) | The CaLP Act defines requirements to: Avoid land degradation Conserve soil Protect water resources Eradicate and prevent the spread and establishment of | Recommend mitigation measures to control pest plant and animal species and to prevent increase in the population of the species as a result of the project. |
| | Enablished and prevent the spread and establishment of noxious weed and pest animal species. This Act defines four categories of noxious weeds: State Prohibited Weeds, Regionally Prohibited Weeds, Regionally Controlled Weeds and Restricted Weeds. Noxious weeds species and the category they are placed in is specific to individual CMA regions Determine whether pest plant or animal species are present within the project area. | |

3. Desktop Result


The following section reviews desktop biodiversity data sourced from relevant Commonwealth and State environmental agencies relevant to the proposed site.

3.1 Native Vegetation

The site occurs in a highly fragmented and modified environment, where historical land use in the form of agriculture (cropping and grazing) and more recent urban/industrial development have significantly altered the landscape.

Figure 3.1 is a modelled vegetation cover produced in 2010. It shows native vegetation persisted on site at this time.

Figure 3.1: Native vegetation extent (DELWP, 2010)



3.1.1 Ecological Vegetation Class (EVC)

The type of native vegetation can be determined from DELWPs Ecological Vegetation Class (EVC) modelling (DSE, 2005). The native vegetation modelled as occurring above, is mapped as Plains Grassland. All other EVCs in a 5km radius can be seen as being remote from the site. DELWP prepared EVC mapping for Victoria in 2005. Figure 3.2 below shows the EVC modelled to occur within and around the site with the location of EVCs discussed in Table 3.1.

Jacobs

Figure 3.2: EVC modelling




Table 3.1: EVCs mapped within 5 km of the project site (DELWP, 2005)

| EVC No. | EVC Name | Bioregion | Bioregional Conservation Status | General Location | Extent of Impact |
|------------|--|-----------|---------------------------------------|--|---|
| 68 | Creekline Grassy Woodland | VVP | Endangered | Associated with the fresh water environment of Hovell's Creek to the north east. | |
| 132 | Plains Grassland | VVP | Endangered | Small isolated patches indicated across the local area. | A review of the endorsed PSP and NVPP indicate no impact |
| 175 | Grassy Woodland | VVP | Endangered | Associated with retained native vegetation in parkland south of the Geelong Ring Road. | to native vegetation on site and no approval requirements relate to the |
| 691 | Aquatic Herbland / Plains Sedgy wetland Mosaic | VVP | Endangered | Associated with the brackish to fresh water environment of Hovell's Creek to the north east. | property. |

3.1.2 Listed Ecological Communities (FFG Act)

The FFG Act lists a number of threatened ecological communities throughout Victoria that are considered to be rare and/or threatened. Table 3.2 provides detail on FFG communities that are considered relevant to the study area. It is noted that the FFG Act currently does not apply to private land, so these risks only apply to road reservations or any public land in the vicinity of the site.

| Listed FFG Community | Analogue | Likely Occurrence | Potential Impact | | |
|--|--|---|--|--|--|
| Western (Basalt) Plains Grasslands Community | EVC 132 Plains Grassland | DELWP mapping has the analogous EVC occurring over a large portion of the site. May occur within road reserves and roadsides in eastern portion of the alignment | None as no native vegetation remains on project site | | |
| Western Basalt Plains (River Red Gum) Grassy Woodland | EVC 55 Plains Grassy Woodland and derived grasslands | Small remnants appear to occur south of the Geelong Ring Road. This community transitions into the grassland community. | None as no native vegetation remains on project site | | |

Table 3.2: FFG threatened ecological communities likely to occur in the area

3.1.3 Threatened Ecological Communities (EPBC)

Threatened Ecological Communities listed under the EPBC Act are modelled to occur in the vicinity of site. The following table (Table 3.3) below.

| EPBC Threatened Ecological Community | Status (EPBC Act) | Modelled Nature of Occurrence | Analogue | Likelihood of Occurrence | Likelihood of Significant Impact |
|---|--------------------------|--|--|--|---|
| Grassy Eucalypt Woodland of the Victorian Volcanic Plain | Critically Endangered | Community known to occur within area | EVC 55 Plains Grassy Woodland | Small remnants appear to occur south of the Geelong Ring Road and remote from site. | Low – TEC unlikely to be present |
| Natural Temperate Grassland of the Victorian Volcanic Plain | Critically Endangered | Community may occur within area | EVC 55 Plains Grassy Woodland and derived grasslands | May occur within road reserves and roadsides with some large patches previously mapped within the site. | Low – TEC unlikely to be present |
| Seasonal Herbaceous Wetlands (Freshwater) of the Temperate Lowland Plains | Critically Endangered | Community likely to occur within area | Several seasonal wetland EVC relate to the EPBC listed community | Cryptic community may occur in shallow depressions in the vicinity of the site. Community difficult to confirm outside of late winter inundation events. | Low - Possibly occurs in and around site in seasonally damp areas. |
| Subtropical and Temperate Coastal Saltmarsh | Vulnerable | Community likely to occur within area | EVC 9 Coastal Saltmarsh | Community does not occur away from tidal marine/estuarine influences. Community likely present on Hovell's Creek to the north east. | Low - Unlikely to occur with the general vicinity of the site. |
| White Box- Yellow Box- Blakely's Red Gum Grassy Woodland and Derived Native Grassland | Critically Endangered | Community likely to occur within area | EVC 55 Plains Grassy Woodland EVC 175 Grassy Woodland | Out of range, community occurs north of the Great Divide. | Low - Unlikely to occur with the general vicinity of the site. |

Table 3.3: Threatened ecological communities modelled to occur in vicinity (5km) of site.

3.1.4 Native Vegetation Preprint Plan

Ecology Partners developed a Native Vegetation Precinct Plan for the Geelong Ring Road Employment Precinct (Ecology Partner's, 2011), mapping remaining native vegetation on all properties within the PSP area.

No native vegetation was mapped within the project site, 164-200 McManus Road, Lara.

3.1.5 Habitat

Site vegetation attributes and other features (wetland, rocky areas) have an important bearing on the habitat offered in a particular area. Figure 3.3 below shows the expected biodiversity values associated with the vicinity of the project site. Green indicates high scores grading to through yellow to brown for low scores. The desktop data suggests the site to have had high biodiversity values, as further considered through the site assessment below.

Figure 3.3: Strategic biodiversity score (DEWLP, 2020)



3.1.6 Site Assessment

A brief site assessment was undertaken to review habitat present on site against desktop data collated. The assessment confirmed the site was highly disturbed and modified through recent site works involving the clearing and levelling of the site and removal of vegetation present. The site has subsequently been used as a refuse dumping ground with dry fill (dirt, bitumen, gravel, rock) deposited across much of the site. Remaining vegetation is largely dominated by weed species.

3.2 Threatened Species

A number of threatened species records have previously been recorded within a 10 km buffer of the site. Although forty-three species have previously been recorded, it is generally considered a low number given the landscape, and likely reflects the limited number of surveys previously undertaken and their results fed into the Victorian Biodiversity Atlas. Table 3.4 provides a summary of the species recorded, while a desktop assessment was undertaken of the records in Table 3.4.

Table 3.4: Total threatened species recorded

| Life Form | Total Recorded | FFG Act | EPBC Act |
|-----------|----------------|---------|----------|
| Birds | 37 | 19 | 9 |
| Insects | 2 | 2 | 1 |
| Mammals | 2 | 1 | 1 |
| Plants | 16 | 5 | 5 |
| Reptiles | 2 | 1 | 1 |
| Total | 59 | 28 | 17 |

Table 3.5: Threatened flora previously recorded (DELWP, 2020)

| Scientific Name | Common Name | Status | Habitat | Records | Last Recorded | Likelihood of Occurrence | Likelihood of a Significant Impact |
|---|--------------------------|--|--|---------|------------------|---|--|
| Atriplex paludosa subsp. paludosa | Marsh Saltbush | VicAdv. Rare | Locally common on the fringes of coastal and near coastal saltmarshes west from Wilsons Promontory. (Walsh and Entwisle 1996) | 1 | 1980 | Low – estuarine species possible in local saline areas. Nearest estuarine environment 4 km to the SE. | Low – Species unlikely to occur in the local area. |
| Avicennia marina subsp. australasica | Grey Mangrove | VicAdv. Rare | Locally common on tidal mudflats, (bays, estuaries and creek mouths) from the western half of Port Phillip Bay to corner inlet. (Gray and Knight 2001) | 3 | 2003 | Low – estuarine species possible in local saline areas. Nearest estuarine environment 4 km to the SE. | Low – Relevant habitat to support species not apparent in close proximity to the site. |
| Comesperma polygaloides | Small Milkwort | FFG VicAdv. Vulnerable | Occasional on heavier soils (clays, alluvium) supporting grassland and grassy woodland communities in central and south western areas. (Walsh and Entwisle 1999) | 1 | 1923 | Low – Species unlikely to persist in local grassland remnants given lack of required ecological processes, particularly fire. | Low – Species unlikely to persist in the local area. |
| Diuris basaltica | Small Golden Moths | EPBC Endangered FFG VicAdv. Endangered | Plains Grassland on Victorian Basalt Plains. Known from records in Laverton and Altona. Flowers SepOct. (Gray and Knight 2001) | 1 | 1998 | Low – Species unlikely to persist in local grassland remnants given lack of required ecological processes, particularly fire. | Low – Species unlikely to persist in the local area. |
| Eucalyptus leucoxylon subsp. connata | Melbourne Yellow-gum | VicAdv. Vulnerable | The main concentration of E. leucoxylon subsp. connata is in the Brisbane Ranges between Bacchus March and Geelong, where it grows on skeletal soils. Also grows on skeletal soils at Long Forest between Bacchus Marsh and Melton, and at Studley Park at Kew (in Melbourne) where it grows on soil derived from Silurian sandstone (RBGV 2018). | 2 | 2011 | Low – No trees are evident in the review of aerial imagery. Specific soil environment required not apparent in the area. | Low – Species unlikely to occur in the local area. |
| Lachnagrostis robusta | Salt Blown- grass | VicAdv. Rare | Occurs around margins of salt lakes and saline depressions mostly across the Volcanic Plain, with eastern outliers near Tooradin and Seaspray and a few sites west of the Grampians (Douglas, Natimuk areas) (RBGV 2016). | 1 | 1997 | Low – estuarine species possible in local saline areas. Nearest estuarine environment 4 km to the SE. | Low – Relevant habitat to support species not apparent in close proximity to the site. |
| Maireana aphylla | Leafless Bluebush | VicAdv. | Rare, occurring on heavy cracking clays and alluvium associated with flood prone sites, considered locally extinct in areas south of | 3 | 2018 | Low – Unlikely to remain in the study area given widescale habitat | Low – Species unlikely to persist in the local area. |

| Scientific Name | Common Name | Status | Habitat | Records | Last Recorded | Likelihood of Occurrence | Likelihood of a Significant Impact |
|---|----------------------------|--|--|---------|------------------|--|---|
| | | Poorly Known | Melbourne, existing records from the north of the state. (Walsh and Entwisle 1996) | | | modification associated with recent development. | |
| Pimelea spinescens subsp. spinescens | Spiny Rice- flower | EPBC Critically Endangered FFG VicAdv. Endangered | Grows in grassland, open shrubland and occasionally woodland, often on basalt-derived soils. Mostly west of Melbourne (to near Horsham), but extending as far north as Echuca (RBGV 2017). | 2 | 2013 | Low - Unlikely to remain in the study area given widescale habitat modification associated with recent development. | Low - Species unlikely to persist in the local area. |
| Podolepis linearifolia | Basalt Podolepis | VicAdv. Endangered | Usually grows on heavy clay soils in grasslands but also recorded for grassy woodlands, open forests and around swamps. Flowers Sep.–Dec. (RBGV 2018) | 1 | 1923 | Low - Unlikely to remain in the study area given widescale habitat modification associated with recent development. | Low – Species unlikely to persist in the local area. |
| Potamogeton australiensis | Thin Pondweed | VicAdv. Poorly Known | Occurs in still or slowing flowing fresh water, in heathland swamps and creeks, on muddy substrate (Walsh and Entwisle 1994). | 1 | 1923 | Low – No aquatic habitat present in close vicinity of site. | Low – Relevant habitat to support species not apparent in close proximity to the site. |
| Prasophyllum spicatum | Dense Leek- orchid | EPBC Vulnerable VicAdv. Endangered | Localised across southern Victoria in coastal heathland and near-coastal heathy forest on sandy soils (RBGV 2015). | 6 | 1934 | Low - Coastal species possible in areas with a greater marine influence. | Low – Relevant habitat to support species not apparent in close proximity to the site. |
| Rutidosis leptorhynchoides | Button Wrinklewort | EPBC Endangered FFG VicAdv. Endangered | In Victoria confined to basaltic grasslands between Rokewood and Melbourne where endangered due to loss of habitat (formerly occurring as far west as Casterton, and on the Gippsland Plain near Newry) (RBGV 2015). | 1 | 1923 | Low - Species habitat not apparent in the immediate vicinity of the site. | Low – Relevant habitat to support species not apparent in close proximity to the site. |
| Rytidosperma richardsonii | Straw Wallaby- grass | VicAdv. Vulnerable | Recorded from grassy woodlands in a few localities in north-east Victoria (e.g. Barnawartha, Dookie, Rutherglen, Springhurst and Yarrawonga). The type specimen of this species was from a plant propagated from wild-collected seed purportedly from Werribee (RBGV 2019). | 1 | 1961 | Low – Evidently the species may have occurred in the area but is no longer considered relevant to the site. | Low – Species unlikely to persist in the local area. |

| Scientific Name | Common Name | Status | Habitat | Records | Last Recorded | Likelihood of Occurrence | Likelihood of a Significant Impact |
|--------------------------|------------------------------|--|---|---------|------------------|---|---|
| Senecio macrocarpus | Large- headed Fireweed | EPBC Vulnerable FFG VicAdv. Endangered | In Victoria largely confined to remnant Themeda grasslands on loamy clay soils derived from basalt from near Melbourne west to Skipton area. Also known from auriferous ground near Stawell. Formerly recorded from near Horsham and Casterton, but apparently long extinct from these areas (Walsh and Entwisle 1999). | 1 | 1998 | Moderate – Species may persist in small degraded grassland remnants on roadsides and potentially the site periphery. | Low – Recent site modification would have removed the species if it persisted in the vicinity of the site. |
| Thelymitra gregaria | Basalt Sun- orchid | FFG VicAdv. Endangered | Endemic to Victoria where found in tussock grassland on red-brown loams derived from basalt (RBGV 2015) | 1 | 1925 | Low – Species unlikely to persist in local grassland remnants given lack of required ecological processes, particularly fire. | Low – Species unlikely to persist in the local area. |
| Tripogonella loliiformis | Rye Beetle- grass | VicAdv. Rare | An uncommon grass of scattered occurrence throughout the state, including rocky areas and the Basalt Plain. (Walsh and Entwisle 1994) | 2 | 1998 | Low - Species unlikely to persist in local grassland remnants given lack of required ecological processes, particularly fire. | Low – Species unlikely to persist in the local area. |

Table 3.6: Threatened fauna previously recorded (DELWP, 2020)

| Scientific Name | Common Name | Status | Habitat | Records | Last Recorded | Likelihood of Occurrence | Likelihood of a Significant Impact | | |
|-----------------------|---------------------|--------------------------------------|---|---------|------------------|--|--|--|--|
| Birds | Birds | | | | | | | | |
| Actitis hypoleucos | Common Sandpiper | VicAdv. Vulnerable | Shallow, pebbly, muddy or sandy edges of rivers and streams, coastal to far inland; dams, lakes, sewage ponds; margins of tidal rivers; waterways in mangroves or saltmarsh; mudflats; rocky or sandy beaches; causeways, riverside lawns, drains, street gutters (Pizzey and Knight 2012) | 1 | 1990 | Low – Species generally associated with aquatic or estuarine habitat. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. | | |
| Anseranas semipalmata | Magpie Goose | FFG VicAdv. Near Threatened | Large seasonal wetlands and well-vegetated dams with rushes and sedges, wet grasslands, floodplains. (Pizzey and Knight 2012) | 8 | 2018 | Low – Wetland species, no wetlands occur within the immediate vicinity of the site. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. | | |

| Scientific Name | Common Name | Status | Habitat | Records | Last Recorded | Likelihood of Occurrence | Likelihood of a Significant Impact |
|-------------------------------|-------------------------|--|---|---------|------------------|---|--|
| Ardea alba | Great Egret | FFG VicAdv. Vulnerable | Shallows of rivers, estuaries, tidal mudflats, freshwater wetlands; sewage ponds, irrigation areas, larger dams etc. (Pizzey and Knight 2012) | 108 | 2019 | Low - Wetland species, no wetlands occur within the immediate vicinity of the site. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Ardea intermedia plumifera | Plumed Egret | FFG VicAdv. Endangered | Freshwater wetlands, pastures and croplands, tidal mudflats, floodplains. (Pizzey and Knight 2012) | 1 | 1977 | Low - Species generally associated with aquatic or estuarine habitat. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Arenaria interpres | Ruddy Turnstone | VicAdv. Vulnerable | Ocean coasts with exposed rock, stony, sandy or shelly beaches, mudflats. Occasionally inland on shallow waters, sewage ponds, saltfields, open or ploughed ground. (Pizzey and Knight 2012) | 1 | 2016 | Low - Species generally associated with aquatic or estuarine habitat. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Aythya australis | Hardhead | VicAdv. Vulnerable | Deep, permanent wetlands, large open waters, brackish coastal swamps, farm dams, ornamental lakes , sewage ponds. (Pizzey and Knight 2012) | 28 | 2018 | Low - Wetland species, no wetlands occur within the immediate vicinity of the site. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Biziura lobata | Musk Duck | VicAdv. Vulnerable | Well-vegetated swamps, wetlands, both brackish and fresh, lakes, reservoirs, shallow bays, inlets; occasionally at sea. (Pizzey and Knight 2012) | 13 | 2018 | Low - Wetland species, no wetlands occur within the immediate vicinity of the site. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Botaurus poiciloptilus | Australasian Bittern | EPBC Endangered FFG VicAdv. Endangered | Narrow habitat preferences, preferring shallow, vegetated freshwater or brackish swamps. (Pizzey and Knight 2012) | 1 | 1990 | Low - Wetland species, no wetlands occur within the immediate vicinity of the site. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Calidris canutus | Red Knot | EPBC Endangered VicAdv. | Tidal mudflats, sandflats, beaches, saltmarshes, flooded pastures, ploughed lands. (Pizzey and Knight 2012) | 1 | 1981 | Low - Species generally associated with aquatic or estuarine habitat. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |

| Scientific Name | Common Name | Status | Habitat | Records | Last Recorded | Likelihood of Occurrence | Likelihood of a Significant Impact |
|------------------------|--------------------------------|--|--|---------|------------------|---|--|
| | | Endangered | | | | | |
| Calidris ferruginea | Curlew Sandpiper | EPBC Critically Endangered FFG VicAdv. Endangered | Tidal mudlfats; saltmarsh, saltfields; fresh, brackish or saline wetlands; sewage ponds. (Pizzey and Knight 2012) | 2 | 2017 | Low - Species generally associated with aquatic or estuarine habitat. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Ceyx azureus | Azure Kingfisher | VicAdv. Near Threatened | Root-festooned banks of fresh or tidal creeks, rivers and streams in rainforest, lakes, swamps, estuaries, mangroves. (Pizzey and Knight 2012) | 1 | 1981 | Low – Generally associated with waterways, may range out over adjoining areas. Site remote from water ways. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Chlidonias hybrida | Whiskered Tern | VicAdv. Near Threatened | Vegetated and open wetlands; brackish, saline lakes; saltfields, irrigated lands, sewage ponds; occasionally offshore. (Pizzey and Knight 2012) | 27 | 2018 | Low - Species generally associated with aquatic or estuarine habitat. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Chlidonias leucopterus | White- winged Black Tern | VicAdv. Near Threatened | large wetlands, coastal and inland; saltfields, sewage ponds, estuaries and coastal waters. (Pizzey and Knight 2012) | 1 | 1977 | Low - Species generally associated with aquatic or estuarine habitat. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Circus assimilis | Spotted Harrier | VicAdv. Near Threatened | Grassy plains, crops and stubblefields; bluebush, saltbush, spinifex associations; scrublands, mallee, heathlands; open, grassy woodlands. (Pizzey and Knight 2012) | 29 | 2019 | High – Species likely to hunt over the study site, with large home ranges associated with the predator. | Low – Specific nesting/breeding habitat not present. |
| Climacteris picumnus | Brown Treecreeper | VicAdv. Near Threatened | Drier forests/woodlands/scrubs, with fallen branches; particularly River Red Gum lined water courses. (Pizzey and Knight 2012) | 1 | 1969 | Low – Associated with woodland habitats which are not present in the vicinity of the site. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |

| Scientific Name | Common Name | Status | Habitat | Records | Last Recorded | Likelihood of Occurrence | Likelihood of a Significant Impact |
|--------------------------|----------------------------------|--|--|---------|------------------|---|--|
| Dromaius novaehollandiae | Emu | VicAdv. Near Threatened | Found in plains, scrublands, open woodlands, coastal heaths, alpine pastures, semi-deserts, margins of lakes, pastoral and cereal growing areas. Mostly absent from closely settled parts, common in pastoral and cropping regions, state forests and national parks (Pizzey and Knight 2012). | 8 | 2019 | Low – Local area becoming increasingly fragmented with recent development. | Low – |
| Egretta garzetta | Little Egret | EPBC Endangered VicAdv. Endangered | Tidal mudflats, saltmarshes, mangroves, freshwater wetlands, sewage ponds. (Pizzey and Knight 2012) | 86 | 2019 | Low - Wetland species, no wetlands occur within the immediate vicinity of the site. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Falco subniger | Black Falcon | FFG VicAdv. Vulnerable | Plains, grasslands, foothills, timbered watercourses, wetland environs; crops; occasionally over towns and cities. (Pizzey and Knight 2012) | 8 | 2018 | High – Species likely to hunt over the study site, with large home ranges associated with the predator. | Low - Specific nesting/breeding habitat not present. |
| Gallinago hardwickii | Latham's Snipe | VicAdv. Near Threatened | Freshwater or brackish wetlands, preferring to be close to protective vegetation cover. (Pizzey and Knight 2012) | 75 | 2019 | Low - Wetland species, no wetlands occur within the immediate vicinity of the site. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Haliaeetus leucogaster | White- bellied Sea- Eagle | FFG VicAdv. Vulnerable | Coasts, inlands, estuaries, inlets, large rivers, inland lakes, reservoirs. (Pizzey and Knight 2012) | 3 | 2018 | High – Species likely to hunt over the study site, with large home ranges associated with the predator. | Low - Specific nesting/breeding habitat not present. |
| Hirundapus caudacutus | White- throated Needletail | EPBC Vulnerable FFG VicAdv. Vulnerable | Airspace over forests, woodlands, farmlands, plains, lakes, coasts, towns, feeding companies frequency patrol back and forward along favoured hilltops and timbered ranges. (Pizzey and Knight 2012) | 3 | 1981 | Moderate – Species likely to over fly the site, however, unlikely to make land fall. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |

| Scientific Name | Common Name | Status | Habitat | Records | Last Recorded | Likelihood of Occurrence | Likelihood of a Significant Impact |
|-----------------------|------------------------------|--|--|---------|------------------|---|--|
| Hydroprogne caspia | Caspian Tern | FFG VicAdv. Near Threatened | Coastal, offshore waters, beaches, mudflats, estuaries, larger rivers, reservoirs and lakes. (Pizzey and Knight 2012) | 13 | 2018 | Low - Species generally associated with aquatic or estuarine habitat. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Ixobrychus dubius | Australian Little Bittern | FFG VicAdv. Endangered | Dense reedbeds in freshwater swamps, lakes and rivers; tussocks in wetland areas. (Pizzey and Knight 2012) | 1 | 1970 | Low - Wetland species, no wetlands occur within the immediate vicinity of the site. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Larus pacificus | Pacific Gull | VicAdv. Near Threatened | Coasts, bays, offshore islands, coastal farmland, swamps, garbage tips; some follow rivers inland. (Pizzey and Knight 2012) | 86 | 2019 | Low - Species generally associated with aquatic or estuarine habitat. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Lathamus discolor | Swift Parrot | EPBC Critically Endangered FFG VicAdv. Endangered | Open grassy woodland, with dead trees, near permanent water and forested hills, coastal heaths, pastures with exotic grasses, weeds, roadsides, orchards. (Pizzey and Knight 2012) | 1 | 1998 | Low – Species annual migration to the Box Ironbark forests to the north may result in the species flying through the area. There are no habitat features the species is likely to utilise in the area. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Lewinia pectoralis | Lewin's Rail | VicAdv. Vulnerable FFG | Swamp woodlands, rushes, reeds, rank grass in swamps, creeks, paddocks; wet heaths. (Pizzey and Knight 2012) | 1 | 1990 | Low - Wetland species, no wetlands occur within the immediate vicinity of the site. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Neophema chrysogaster | Orange- bellied Parrot | EPBC Critically Endangered FFG VicAdv. | On mainland prefers small islands, peninsulas in coastal areas; with saltmarsh plants; coastal pastures, golf courses, crops of millet and sunflowers; dunes, beaches. Tasmania prefers button-grass; sedges on wet peat plains and eucalypt woodland on margin (Pizzey and Knight 2012) | 1 | 1977 | Low - Species generally associated with aquatic or estuarine habitat. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |

| Scientific Name | Common Name | Status | Habitat | Records | Last Recorded | Likelihood of Occurrence | Likelihood of a Significant Impact |
|------------------------------|--------------------------|--|--|---------|------------------|---|--|
| | | Critically Endangered | | | | | |
| Ninox connivens | Barking Owl | FFG VicAdv. Endangered | Open forests, woodlands, dense scrubs, foothills, river red gums, other large trees near water courses, penetrating otherwise open country, and paperbark woodlands. (Pizzey and Knight 2012) | 1 | 1969 | High – Species likely to hunt over the study site, with large home ranges associated with the predator. | Low - Specific nesting/breeding habitat not present. |
| Ninox strenua | Powerful Owl | FFG VicAdv. Vulnerable | Pairs occupy a large, probably permanent, home range in mountain forests, gullies and forest margins, sparser hilly woodlands, coastal forests, woodlands, scrubs, exotic pine plantations, large trees in private/public gardens, some in cities. (Pizzey and Knight 2012) | 1 | 1969 | High – Species likely to hunt over the study site, with large home ranges associated with the predator. | Low - Specific nesting/breeding habitat not present. |
| Numenius madagascariensis | Eastern Curlew | EPBC Critically Endangered FFG VicAdv. Vulnerable | Estuaries, tidal mudflats, sandspits, saltmarshes, mangroves; occasionally fresh or brackish lakes; bare grasslands near water. (Pizzey and Knight 2012) | 3 | 1992 | Low - Species generally associated with aquatic or estuarine habitat. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Nycticorax caledonicus | Nankeen Night-Heron | VicAdv. Near Threatened | Shallow margins of rivers, wetlands, mangrove- lined estuaries, offshore islands, floodwaters, garden trees. (Pizzey and Knight 2012) | 5 | 1999 | Low - Wetland species, no wetlands occur within the immediate vicinity of the site. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Oxyura australis | Blue-billed Duck | FFG VicAdv. Endangered | Found on temperate, fresh to saline, terrestrial wetlands including sewerage ponds, rivers, salt lakes and saltpans. Preferring deep, permanent open water within or near dense vegetation. (Pizzey and Knight 2012) | 2 | 2013 | Low - Wetland species, no wetlands occur within the immediate vicinity of the site. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Phalacrocorax fuscescens | Black-faced Cormorant | VicAdv. Near Threatened | Coastal waters: islands, larger bays, deep inlets; ashore mostly on isolated rock-stacks, remote rocky headlands; seldom on beaches. (Pizzey and Knight 2012) | 3 | 2019 | Low - Species generally associated with aquatic or estuarine habitat. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |

| Scientific Name | Common Name | Status | Habitat | Records | Last Recorded | Likelihood of Occurrence | Likelihood of a Significant Impact |
|----------------------|--------------------------|--|---|---------|------------------|---|--|
| Phalacrocorax varius | Pied Cormorant | VicAdv. Near Threatened | Coastal waters with sloping shorelines; estuaries, bays, tidal inlets, large inland lakes and rivers, irrigation ponds, coastal mangroves and offshore islands. (Pizzey and Knight 2012) | 98 | 2019 | Low - Species generally associated with aquatic or estuarine habitat. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Platalea regia | Royal Spoonbill | VicAdv. Near Threatened | Larger shallow waters, inland and coastal, well- vegetated shallow freshwater wetlands, saltfields, mangroves, islands, farm dams occasionally. (Pizzey and Knight 2012) | 101 | 2019 | Low - Species generally associated with aquatic or estuarine habitat. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Plegadis falcinellus | Glossy Ibis | VicAdv. Near Threatened | Well vegetated wetlands, wet pastures, rice fields, floodwaters, floodplains, brackish or occasionally saline wetlands, mangroves, mudflats; occasionally dry grasslands. (Pizzey and Knight 2012) | 1 | 1991 | Low - Wetland species, no wetlands occur within the immediate vicinity of the site. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Spatula rhynchotis | Australasian Shoveler | VicAdv. Vulnerable | Larger waters, fresh and saline lakes, well- vegetated freshwater wetlands, coastal inlets, sewage ponds, floodwaters. (Pizzey and Knight 2012) | 2 | 1991 | Low - Wetland species, no wetlands occur within the immediate vicinity of the site. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Sternula albifrons | Little Tern | FFG VicAdv. Vulnerable | Coastal waters, bays, inlets, saline or brackish lakes, saltfields, sewerage ponds near coast. (Pizzey and Knight 2012) | 7 | 2018 | Low - Species generally associated with aquatic or estuarine habitat. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Sternula nereis | Fairy Tern | EPBC Vulnerable FFG VicAdv. Endangered | Coastal waters, bays, inlets, saline or brackish lakes, saltfields, sewerage ponds near coast. Breeds Sept-Jan in single pairs to large colonies on beaches, islands, rock platforms from north of Broome to eastern Victoria/NSW border; much lesser numbers in south (Pizzey and Knight 2012) | 11 | 2018 | Low - Species generally associated with aquatic or estuarine habitat. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Stictonetta naevosa | Freckled Duck | FFG VicAdv. Endangered | Large, well vegetated swamps; in dry periods moves to open lakes. (Pizzey and Knight 2012) | 8 | 2014 | Low - Wetland species, no wetlands occur within the immediate vicinity of the site. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |

| Scientific Name | Common Name | Status | Habitat | Records | Last Recorded | Likelihood of Occurrence | Likelihood of a Significant Impact |
|---------------------------|--|--|---|---------|------------------|--|---|
| Tringa nebularia | Common Greenshank | VicAdv. Vulnerable | Mudflats, estuaries, saltmarshes, margins of lakes; wetlands, claypans, fresh and saline; commercial saltfields and sewage ponds. (Pizzey and Knight 2012) | 6 | 2014 | Low - Species generally associated with aquatic or estuarine habitat. | Low – Suitable habitat not present in the immediate vicinity of site, therefore the species is unlikely to be present. |
| Insects | | | | | | | |
| Hesperilla flavescens | Yellow Sedge- skipper Butterfly | FFG VicAdv. Vulnerable | Poorly understood species thought to be associated with coastal areas containing stands of Gahnia filum (Coleman, 2000) | 1 | 1988 | Low – Relevant habitat not present in the wider vicinity of the site. | Low – Species unlikely to be present. |
| Synemon plana | Golden Sun Moth | EPBC Critically Endangered FFG VicAdv. Critically Endangered | Native temperate grassland and open grassy woodlands, may also be found in degraded grasslands dominated by exotic Chilean Needlegrass (DAWE 2020a) | 2 | 2017 | Moderate – Local area subject to habitat loss, with recent development removing specific tussocked grassland habitat. | Low – the small areas of remaining tussocked grass is composed of exotic species and the extent remaining is considered too small to support a population. |
| Mammals | I | | | I | 1 | I | |
| Perameles gunnii | Eastern Barred Bandicoot | FFG EPBC Vulnerable | Extinct in the wild and occurs in small reintroduced populations in southern Victoria. Previously it occurred across the Victorian Volcanic Plains. During the day it rests in a grass nest. It feeds on earthworms, insects, bulbs, tubers and fungi throughout the night (DAWE 2020b) | 15 | 1977 | Low – Local area becoming increasingly fragmented with recent development limiting the chances of the species persisting. Species assumed to be locally extinct. | Low – Species unlikely to be present. |
| Sminthopsis crassicaudata | Fat-tailed Dunnart | VicAdv. Near Threatened | Open woodland, low shrublands of saltbush and bluebush, tussock grasslands on clay or sandy soils, gibber plain and farmlands. (Van Dyck and Strahan 2008) | 1 | 1972 | Low – Local area becoming increasingly fragmented with recent development. | Low – Species unlikely to be present. |

Qualified Desktop Flora and Fauna Assessment



| Scientific Name | Common Name | Status | Habitat | Records | Last Recorded | Likelihood of Occurrence | Likelihood of a Significant Impact |
|--------------------------|------------------------------|--|--|---------|------------------|---|--|
| Reptiles | Reptiles | | | | | | |
| Delma impar | Striped Legless Lizard | EPBC Vulnerable FFG VicAdv. Vulnerable | Intact native grassland areas, usually in or around stony rise country or deeply cracked earth. (Cogger 2014) | 1 | 1992 | Moderate – While the local area becoming increasingly fragmented with recent development removing much of the specific tussocked grassland habitat, the cryptic species may remain in exotic grassland habitat. | Low – Species unlikely to be present. |
| Pseudemoia pagenstecheri | Tussock Skink | VicAdv. Vulnerable | Tussock grasslands with few or no trees from highlands in ne Victoria to low-altitude basalt plains of sthn Vic (Wilson and Swan 2008) | 2 | 2017 | Low – Local area becoming increasingly fragmented with recent development removing specific tussocked grassland habitat. | Low – Species unlikely to be present. |

3.2.1 Threatened Species (PMST)

The Protected Maters Search Tool (PMST) models Commonwealth listed threatened species and communities within the 5 km search area. Given the modelled nature of the result, many species related to marine (Port Philip, Limeburners Bay and Corio Bay) are included in the result, picking up a number of albatross, turtle, whale and shark species. These are obviously not relevant to the site and are given a 'Low' likelihood of occurrence accordingly. Many other species are grassland specialists that have been heavily impacted by land use change (hence listing) and have limited potential to remain in the local area and specifically the site, given the lack of habitat present. There are however a small number of listed species that have a cryptic ecology and may remain in the modified landscape as discussed below.

The PMST results are assessed in Appendix A, with most of the sixty-one resulting species considered to have a Low likelihood of occurrence. Seven species are considered to have a moderate likelihood of occurrence in the immediate vicinity of the site and are listed in Table 3.7 below.

| Species | EPBC Status | Rationale | Likelihood of a Significant Impact |
|---|--------------------------|---|--|
| Amphibian | | | |
| <i>Litoria raniformis</i> Growling Grass Frog | Vulnerable | Species occurs in and around drainage lines and seasonally wet areas. The site retains some connectivity to such environments, particularly in the north western corner of the site. Species has not been recorded in local area. | Unlikely given site modification and disturbance. Limited habitat observed on site. |
| Invertebrate | | | |
| <i>Synemon plana</i> Golden Sun Moth | Critically Endangered | Highly cryptic species that may persist or establish in areas of degraded and/or exotic grassland retaining <i>Rytidosperma</i> and/or <i>Nassella</i> species. Tussocked grassland areas are evident within the aerial photography. | Unlikely given site modification and disturbance. Limited habitat observed on site. |
| Plants | | | |
| <i>Dianella amoena</i> Matted Flax-lily | Endangered | Species that may persist in otherwise disturbed environments. Species has not been recorded in local area. | Unlikely given site modification and disturbance. |
| Lachnagrostis adamsonii Adamson's Blown- grass | Endangered | Species that may persist in otherwise disturbed environments, particularly where saline drainage concentrates. | Unlikely given site modification and disturbance. |
| Pimelea spinescens subsp. spinescens Plains Rice-flower | Critically Endangered | Species that may persist in otherwise disturbed environments. | Unlikely given site modification and disturbance. |
| Senecio macrocarpus Large-fruit Fireweed | Vulnerable | Species that may persist in otherwise disturbed environments. | Unlikely given site modification and disturbance. |
| Reptiles | | | |

Table 3.7: PMST species considered with moderate likelihood of occurrence

| Species | EPBC Status | Rationale | Likelihood of a Significant Impact |
|---|-------------|---|--|
| <i>Delma impar</i> Striped Legless Lizard | Vulnerable | Highly cryptic species that may persist or establish in areas of degraded and/or exotic grassland retaining <i>Rytidosperma</i> and/or <i>Nassella</i> species. Tussocked grassland areas are evident within the aerial photography. | Unlikely given site modification and disturbance. |

4. Assessment

Based on the Preliminary assessment undertaken, an understanding of potential approval requirements and further assessment works are outlined in Table 4.1. It is important to note that the site has been significantly disturbed and modified following the approval of the Geelong Ring Road Employment Precinct Plan, so results obtained through the review or relevant data bases need to be considered in relation to recent site activities.

Based on the design information currently available and the site attributes identified through this assessment, there are no threatened flora and fauna risks relevant to the site given lack of remaining habitat. Native vegetation was not identified on site during the recent site assessment or recorded in the preparation of the relevant Precinct Structure Plan (Ecology Partners, 2011). The site was found to be largely covered in fill dumped after the vegetation cover was removed and the site levelled, with the majority of surface rock (basalt) outcrops having been removed in recent years. Vegetation present was largely composed of weed species including, but not limited to:

- Serrated Tussock (Nassella trichotoma);
- Toowoomba Canary-grass (Phalaris aquatica);
- Mustard (Brassica spp.);
- Scotch Thistle (Onopordum acanthium);
- Artichoke Thistle (Cynara cardunculus subsp. flavescens);
- Variegated Thistle (Silybum marianum);
- Plantain (Plantago lanceolate);
- Oat (Avena spp.);
- Chilean Needle-grass (Nassella neesiana);
- African Boxthorn (Lycium ferocissimum);
- Couch (Cynodon dactylon var. dactylon);
- Soursob (Oxalis pes-caprae);
- Paterson's curse (Echium plantagineum);
- Galenia (Galenia spp.); and
- Rye grass (Lolium perenne).

Consideration of the potential risks to the project based on the current design are further detailed in Table 4.1 below.

| Policy/legislation | Project Risk | Further Requirement |
|--|--|--|
| Commonwealth | | |
| Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act) | Threatened Ecological Communities Potential for three threatened ecological communities to occur on site. • Natural Temperate Grassland of the Victorian Volcanic Plain • Grassy Woodlands and derived Grasslands of the Victorian Volcanic Plain • Grassy Woodlands and derived Grasslands of the Victorian Volcanic Plain • Seasonal Herbaceous Wetlands (Freshwater) of the Temperate Lowland Plains Vegetation mapping undertaken by DELWP indicate the presence of the synonymous plains grassland EVC across much of the site. The NVPP endorsed by GGCC indicated no native vegetation remains. Threatened Species Potential for three EPBC listed species to occur on the VVP: | Threatened Ecological CommunitiesIt is considered unlikely that the EPBC listedGrassland communities remain on site givenrecent site clearing activities.Threatened SpeciesBoth the Golden Sun Moth and StripedLegless Lizard are known from the area andboth species can inhabit relatively degradedsites as long as the structural tussockelement of grasses (which can be weed)persist. An assessment of specific sitecharacter suggests the small area ofpotential habitat identified through thedesktop assessment (aerial photographyinterpretation) was unlikely to support thegrassland species given the structure andspecies composition of species present. |

Jacobs

| Policy/legislation | Project Risk | Further Requirement | |
|---|---|--|--|
| | Golden Sun MothStriped Legless lizardSpiny Rice Flower | | |
| State | | | |
| Environment Effects Act 1978 (EE Act) | The proposal was considered against relevant EES triggers as they relate to flora and fauna: <u>Native vegetation clearance</u> No native vegetation is reported to occur within the project footprint. Refer Section 3.1 <u>Threatened species</u> The project will not lead to the long-term loss of a significant proportion of known remaining habitat or population of a threatened species within Victoria. Wetlands The site is remote from significant wetlands and Ramsar sites <u>Aquatic, estuarine or marine impacts</u> The project is remote from these systems so will not have_extensive or major effects on the health or biodiversity. | No ecological triggers for an EES are met by the project. | |
| Flora and Fauna Guarantee Act 1988 (FFG Act) | The project does not involve public lands, so the FFG Act is not invoked. | No FFG permits are considered relevant for the project given land tenure. | |
| DELWP Victorian Advisory Lists (VicAdv) | The Victorian Advisory list is considered as part of the threatened species triggers for the EES requirement and as potential specific credits should native vegetation permits be considered for threatened species habitat removal. | No Victorian Advisory listed species are considered relevant to the project. | |
| Planning and Environment Act 1987 (P&E Act) | The site occurs within the Geelong Ring Road Employment Precinct, where a Native Vegetation Precinct Structure Plan | No further assessment or approvals required. | |
| Guidelines for the removal, destruction or lopping of native vegetation (the Guidelines) | has been endorsed by Council under Clause 52.16 of the P&E Act. No native vegetation was identified on site in the NVPP and no retention areas occur on site. | | |
| Catchment and Land Protection Act 1994 (CaLP Act) | Noxious weed management considerations will need to be applied to the project. | Project EMP required to included weed management for protection to meet CaLP Act requirements. | |

5. Conclusion and recommendations

The desktop assessment was undertaken for the proposed Prospect Hill EfW Project at 164-200 McManus Road, Lara.

The data base review found a number of issues pertaining to native vegetation and threatened species on site, however, more recent reporting, particularly that associated with the development of the Geelong Ring Road Precinct Structure Plan, and the accompanying Native Vegetation Precinct Plan had determined that native vegetation did not persist on site. Obvious site development has been undertaken further compromising the likelihood of significant statutory issues relating to flora and fauna.

A brief habitat assessment was undertaken to determine the need for any further targeted threatened species assessments. This found the site to be substantially modified and disturbed and habitat present was considered unsuitable for any of the threatened species considered as potential relevant to the site through the desktop data review,

This report concludes there are no statutory ecological issues relevant to the site.

6. References

Coleman, P. 2000. Local Recovery Plan for the Yellowish sedge-skipper and thatching grass. South Australian Urban Forest Biodiversity Program

Cogger, H. (2014). Reptiles and Amphibians of Australia. Clayton, CSIRO Publishing.

DAWE (2020). "Protected Matters Search Tool." from <u>http://www.environment.gov.au/epbc/protected-matterssearch-tool</u>.

DAWE (2020a). Species Profile and Threats Database: Synemom plana - Golden Sun Moth. Retrieved 9/3/2020, http://www.environment.gov.au/sprats Department of Agriculture, Water and the Environment, Government of Australia

DELWP (2017). Guidelines for the removal, destruction or lopping of native vegetation. Department of Environment, Land, Water and Planning, Government of Victoria. Melbourne.

DELWP (2020). "Victorian Biodiversity Atlas." from http://www.depi.vic.gov.au/environment-andwildlife/ biodiversity/Victorian-biodiversity-atlas. Department of Environment, Land, Water and Planning, Government of Victoria. Melbourne.

DELWP, (2020) Ecological vegetation Class Data. Provided under License. Department of Environment, Water, Land and Planning, Melbourne.

DoE (2016) Threatened Species Scientific Committee - Conservation Advice *Delmar impar* (Striped Legless Lizard). Unpublished report, Australian Government Department of the Environment, Canberra.

DSE (2004). Vegetation quality assessment manual: Guidelines for applying the Habitat Hectares scoring method. Version 1.3. Melbourne, Victorian Department of Sustainability and Environment.

Ecology Partners, 2011 Native Vegetation Precinct Plan, Geelong Ring Road Employment Precinct. City of Greater Geelong.

Gray, M. and J. Knight (2001). Flora of Melbourne: A guide to the indigneous plants of the Greater Melbourne Area. Hyland House, Flemington.

Menkhorst, P. & Broome, L. (2006) National Recovery Plan for the Smoky Mouse Pseudomys fumeus. State of Victoria Department of Sustainability and Environment 2006.

Pizzey, G. and F. Knight (2012). The Field Guide to the Birds of Australia. Sydney, Harper Collins Australia.

RBGV (2015). Flora of Victoria Online (VicFlora). <https://vicflora.rbg.vic.gov.au/> Royal Botanic Gardens Victoria.

RBGV (2016). Flora of Victoria Online (VicFlora). <https://vicflora.rbg.vic.gov.au/> Royal Botanic Gardens Victoria.

RBGV (2017). Flora of Victoria Online (VicFlora). <https://vicflora.rbg.vic.gov.au/> Royal Botanic Gardens Victoria.

RBGV (2018). Flora of Victoria Online (VicFlora). <https://vicflora.rbg.vic.gov.au/> Royal Botanic Gardens Victoria.

RBGV (2019). Flora of Victoria Online (VicFlora). <https://vicflora.rbg.vic.gov.au/> Royal Botanic Gardens Victoria.

Van Dyck, S. and R. Strahan (2008). The Mammals of Australia. Reed New Holland, Sydney.

Walsh, N. and T. Entwisle (1994). Flora of Victoria Volume 2: Ferns and Allied Plants, Conifers and Monocotyledons, Inkata Press, Melbourne.

Walsh, N. and T. Entwisle (1996). Flora of Victoria Volume 3: Dicotoledons; Winteraceae to Mytaceae, Inkata Press, Melbourne.

Walsh, N. and T. Entwisle (1999). Flora of Victoria Volume 4: Dicotoledons; Cornaceae to Asteraceae, Inkata Press, Melbourne.

Wilson, S. and G. Swan (2008). A Complete Guide to Reptiles of Australia - Second Edition. Reed New Holland, Sydney.



Appendix A. Protected Matters Search Tool Report



EPBC Act Protected Matters Report

This report provides general guidance on matters of national environmental significance and other matters protected by the EPBC Act in the area you have selected.

Information on the coverage of this report and qualifications on data supporting this report are contained in the caveat at the end of the report.

Information is available about <u>Environment Assessments</u> and the EPBC Act including significance guidelines, forms and application process details.

Report created: 23/04/20 16:32:10

Summary Details Matters of NES Other Matters Protected by the EPBC Act Extra Information Caveat Acknowledgements



This map may contain data which are ©Commonwealth of Australia (Geoscience Australia), ©PSMA 2010

| <u>Coordinates</u> | |
|--------------------|--|
| Buffer: 5.0Km | |
| | |



Summary

Matters of National Environmental Significance

This part of the report summarises the matters of national environmental significance that may occur in, or may relate to, the area you nominated. Further information is available in the detail part of the report, which can be accessed by scrolling or following the links below. If you are proposing to undertake an activity that may have a significant impact on one or more matters of national environmental significance then you should consider the Administrative Guidelines on Significance.

| World Heritage Properties: | None |
|---|------|
| National Heritage Places: | None |
| Wetlands of International Importance: | 1 |
| Great Barrier Reef Marine Park: | None |
| Commonwealth Marine Area: | None |
| Listed Threatened Ecological Communities: | 5 |
| Listed Threatened Species: | 61 |
| Listed Migratory Species: | 60 |

Other Matters Protected by the EPBC Act

This part of the report summarises other matters protected under the Act that may relate to the area you nominated. Approval may be required for a proposed activity that significantly affects the environment on Commonwealth land, when the action is outside the Commonwealth land, or the environment anywhere when the action is taken on Commonwealth land. Approval may also be required for the Commonwealth or Commonwealth agencies proposing to take an action that is likely to have a significant impact on the environment anywhere.

The EPBC Act protects the environment on Commonwealth land, the environment from the actions taken on Commonwealth land, and the environment from actions taken by Commonwealth agencies. As heritage values of a place are part of the 'environment', these aspects of the EPBC Act protect the Commonwealth Heritage values of a Commonwealth Heritage place. Information on the new heritage laws can be found at http://www.environment.gov.au/heritage

A <u>permit</u> may be required for activities in or on a Commonwealth area that may affect a member of a listed threatened species or ecological community, a member of a listed migratory species, whales and other cetaceans, or a member of a listed marine species.

| None |
|------|
| None |
| 71 |
| 7 |
| None |
| None |
| None |
| |

Extra Information

This part of the report provides information that may also be relevant to the area you have nominated.

| State and Territory Reserves: | 2 |
|----------------------------------|------|
| Regional Forest Agreements: | 1 |
| Invasive Species: | 34 |
| Nationally Important Wetlands: | 1 |
| Key Ecological Features (Marine) | None |

Details

Matters of National Environmental Significance

| Wetlands of International Importance (Ramsar) | [Resource Information] |
|--|------------------------|
| Name | Proximity |
| Port phillip bay (western shoreline) and bellarine peninsula | Within Ramsar site |

| Listed Threatened Ecological Communities | | [Resource Information] | |
|--|-----------------------|--|--|
| For threatened ecological communities where the distribution is well known, maps are derived from recovery plans, State vegetation maps, remote sensing imagery and other sources. Where threatened ecological community distributions are less well known, existing vegetation maps and point location data are used to produce indicative distribution maps. | | | |
| Name | Status | Type of Presence | |
| Grassy Eucalypt Woodland of the Victorian Volcanic Plain | Critically Endangered | Community known to occur within area | |
| Natural Temperate Grassland of the Victorian Volcanic Plain | Critically Endangered | Community likely to occur within area | |
| Seasonal Herbaceous Wetlands (Freshwater) of the Temperate Lowland Plains | Critically Endangered | Community likely to occur within area | |
| Subtropical and Temperate Coastal Saltmarsh | Vulnerable | Community likely to occur within area | |
| White Box-Yellow Box-Blakely's Red Gum Grassy Woodland and Derived Native Grassland | Critically Endangered | Community likely to occur within area | |
| Listed Threatened Species | | [Resource Information] | |
| Name | Status | Type of Presence | |
| Birds | | | |
| Anthochaera phrygia | | | |
| Regent Honeyeater [82338] | Critically Endangered | Foraging, feeding or related behaviour may occur within area | |
| Botaurus poiciloptilus | | | |
| Australasian Bittern [1001] | Endangered | Species or species habitat known to occur within area | |
| Calidris canutus | | | |
| Red Knot, Knot [855] | Endangered | Species or species habitat known to occur within area | |
| Calidris ferruginea | | | |
| Curlew Sandpiper [856] | Critically Endangered | Species or species habitat known to occur within area | |
| Calidris tenuirostris | | | |
| Great Knot [862] | Critically Endangered | Species or species habitat known to occur within area | |
| Charadrius leschenaultii | | | |
| Greater Sand Plover, Large Sand Plover [877] | Vulnerable | Species or species habitat known to occur within area | |
| Charadrius mongolus | | | |
| Lesser Sand Plover, Mongolian Plover [879] | Endangered | Species or species habitat known to occur within area | |
| Diomedea antipodensis | | | |
| Antipodean Albatross [64458] | Vulnerable | Foraging, feeding or related behaviour likely to occur within area | |

| Name | Status | Type of Presence |
|---|-----------------------|--|
| Diomedea epomophora Southern Royal Albatross [89221] Diomedea exulans | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| Wandering Albatross [89223] | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| Diomedea sanfordi Northern Royal Albatross [64456] | Endangered | Foraging, feeding or related behaviour likely to occur within area |
| Grantiella picta Painted Honeyeater [470] | Vulnerable | Species or species habitat known to occur within area |
| Hirundapus caudacutus White-throated Needletail [682] | Vulnerable | Species or species habitat likely to occur within area |
| Lathamus discolor Swift Parrot [744] | Critically Endangered | Species or species habitat known to occur within area |
| Limosa lapponica baueri Bar-tailed Godwit (baueri), Western Alaskan Bar-tailed Godwit [86380] | Vulnerable | Species or species habitat known to occur within area |
| Limosa lapponica menzbieri Northern Siberian Bar-tailed Godwit, Bar-tailed Godwit (menzbieri) [86432] | Critically Endangered | Species or species habitat may occur within area |
| Macronectes giganteus Southern Giant-Petrel, Southern Giant Petrel [1060] | Endangered | Species or species habitat may occur within area |
| Macronectes halli Northern Giant Petrel [1061] | Vulnerable | Species or species habitat may occur within area |
| Neophema chrysogaster Orange-bellied Parrot [747] | Critically Endangered | Migration route likely to occur within area |
| Numenius madagascariensis Eastern Curlew, Far Eastern Curlew [847] | Critically Endangered | Species or species habitat known to occur within area |
| Pachyptila turtur_subantarctica Fairy Prion (southern) [64445] | Vulnerable | Species or species habitat known to occur within area |
| Pedionomus torquatus Plains-wanderer [906] | Critically Endangered | Species or species habitat likely to occur within area |
| Phoebetria fusca Sooty Albatross [1075] | Vulnerable | Species or species habitat likely to occur within area |
| Pterodroma leucoptera leucoptera Gould's Petrel, Australian Gould's Petrel [26033] | Endangered | Species or species habitat may occur within area |
| Rostratula australis Australian Painted Snipe [77037] | Endangered | Species or species habitat known to occur within area |
| <u>Sternula nereis_nereis</u> Australian Fairy Tern [82950] | Vulnerable | Species or species habitat known to occur within area |
| <u>Thalassarche bulleri</u> Buller's Albatross, Pacific Albatross [64460] | Vulnerable | Species or species |

| Name | Status | Type of Presence |
|--|-----------------------|---|
| | | habitat may occur within area |
| Thalassarche bulleri platei | | alea |
| Northern Buller's Albatross, Pacific Albatross [82273] | Vulnerable | Species or species habitat may occur within area |
| | | may occur within area |
| Thalassarche cauta_cauta Shy Albatross [82345] | Vulnerable | Foraging, feeding or related |
| Shy Abalioss [02343] | Vullerable | behaviour likely to occur |
| Thalassarche cauta steadi | | within area |
| White-capped Albatross [82344] | Vulnerable | Foraging, feeding or related |
| | | behaviour likely to occur within area |
| Thalassarche chrysostoma | | |
| Grey-headed Albatross [66491] | Endangered | Species or species habitat may occur within area |
| | | may occur within area |
| Thalassarche impavida | Vulnorabla | Foraging, feeding or related |
| Campbell Albatross, Campbell Black-browed Albatross [64459] | vuinerable | behaviour likely to occur |
| Thalassarche melanophris | | within area |
| Black-browed Albatross [66472] | Vulnerable | Species or species habitat |
| | | may occur within area |
| Thalassarche salvini | | |
| Salvin's Albatross [64463] | Vulnerable | Foraging, feeding or related |
| | | behaviour likely to occur within area |
| Thinornis rubricollis rubricollis | ., | 0 |
| Hooded Plover (eastern) [66726] | Vulnerable | Species or species habitat likely to occur within area |
| Fish | | , |
| Galaxiella pusilla | | |
| Eastern Dwarf Galaxias, Dwarf Galaxias [56790] | Vulnerable | Species or species habitat |
| | | likely to occur within area |
| Prototroctes maraena | | . |
| Australian Grayling [26179] | Vulnerable | Species or species habitat likely to occur within area |
| F | | |
| Frogs Litoria raniformis | | |
| Growling Grass Frog, Southern Bell Frog, Green and | Vulnerable | Species or species habitat |
| Golden Frog, Warty Swamp Frog, Golden Bell Frog [1828] | | known to occur within area |
| Insects | | |
| Synemon plana | Critically Endeparted | Spaciae or operioe hebitet |
| Golden Sun Moth [25234] | Critically Endangered | Species or species habitat known to occur within area |
| Mammals | | |
| Eubalaena australis | | |
| Southern Right Whale [40] | Endangered | Species or species habitat |
| | | known to occur within area |
| Megaptera novaeangliae | | . |
| Humpback Whale [38] | Vulnerable | Species or species habitat may occur within area |
| | | |
| Pteropus poliocephalus Grey-headed Flying-fox [186] | Vulnerable | Foraging, feeding or related |
| | Valiforable | behaviour likely to occur |
| Plants | | within area |
| Amphibromus fluitans | | |
| River Swamp Wallaby-grass, Floating Swamp Wallaby-grass [19215] | Vulnerable | Species or species habitat likely to occur within area |
| vvaliaby-grass [13210] | | INCIY IO OCCUI WILIIII AICA |

| Name | Status | Type of Presence |
|--|-----------------------|---|
| Dianella amoena Matted Flax-lily [64886] | Endangered | Species or species habitat may occur within area |
| Diuris basaltica Small Golden Moths Orchid, Early Golden Moths [64654] | Endangered | Species or species habitat may occur within area |
| Dodonaea procumbens Trailing Hop-bush [12149] | Vulnerable | Species or species habitat may occur within area |
| <u>Glycine latrobeana</u> Clover Glycine, Purple Clover [13910] | Vulnerable | Species or species habitat likely to occur within area |
| <u>Lachnagrostis adamsonii</u> Adamson's Blown-grass, Adamson's Blowngrass [76211] | Endangered | Species or species habitat may occur within area |
| Leucochrysum albicans var. tricolor Hoary Sunray, Grassland Paper-daisy [56204] | Endangered | Species or species habitat likely to occur within area |
| Pimelea spinescens subsp. spinescens Plains Rice-flower, Spiny Rice-flower, Prickly Pimelea [21980] | Critically Endangered | Species or species habitat known to occur within area |
| Prasophyllum frenchii Maroon Leek-orchid, Slaty Leek-orchid, Stout Leek- orchid, French's Leek-orchid, Swamp Leek-orchid [9704] | Endangered | Species or species habitat likely to occur within area |
| Pterostylis cucullata Leafy Greenhood [15459] | Vulnerable | Species or species habitat may occur within area |
| Rutidosis leptorrhynchoides Button Wrinklewort [7384] | Endangered | Species or species habitat likely to occur within area |
| Senecio macrocarpus Large-fruit Fireweed, Large-fruit Groundsel [16333] | Vulnerable | Species or species habitat likely to occur within area |
| Xerochrysum palustre Swamp Everlasting, Swamp Paper Daisy [76215] | Vulnerable | Species or species habitat may occur within area |
| Reptiles | | |
| <u>Caretta caretta</u> Loggerhead Turtle [1763] | Endangered | Species or species habitat known to occur within area |
| <u>Chelonia mydas</u> Green Turtle [1765] | Vulnerable | Foraging, feeding or related behaviour known to occur within area |
| Delma impar Striped Legless Lizard, Striped Snake-lizard [1649] | Vulnerable | Species or species habitat known to occur within area |
| Dermochelys coriacea Leatherback Turtle, Leathery Turtle, Luth [1768] | Endangered | Species or species habitat known to occur within area |
| Tympanocryptis pinguicolla Grassland Earless Dragon [66727] | Endangered | Species or species habitat may occur within area |
| Sharks | | |
| <u>Carcharodon carcharias</u> White Shark, Great White Shark [64470] | Vulnerable | Species or species habitat known to occur |

| Name | Status | Type of Presence |
|--|--------------------------|--|
| | | within area |
| Listed Migratory Species * Species is listed under a different scientific name on th | he EPBC Act - Threatened | [Resource Information] Species list. |
| Name | Threatened | Type of Presence |
| Migratory Marine Birds | | |
| Anous stolidus Common Noddy [825] | | Species or species habitat likely to occur within area |
| Apus pacificus Fork-tailed Swift [678] | | Species or species habitat likely to occur within area |
| Ardenna carneipes Flesh-footed Shearwater, Fleshy-footed Shearwater [82404] | | Foraging, feeding or related behaviour likely to occur within area |
| Ardenna grisea Sooty Shearwater [82651] | | Species or species habitat may occur within area |
| Diomedea antipodensis | | |
| Antipodean Albatross [64458] | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| Diomedea epomophora Southern Royal Albatross [89221] | Vulnerable | Foraging, feeding or related |
| | Vanorabio | behaviour likely to occur within area |
| Diomedea exulans Wandering Albatross [89223] | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| Diomedea sanfordi Northern Royal Albatross [64456] | Endangered | Foraging, feeding or related behaviour likely to occur within area |
| Macronectes giganteus Southern Giant-Petrel, Southern Giant Petrel [1060] | Endangered | Species or species habitat may occur within area |
| Macronectes halli | | |
| Northern Giant Petrel [1061] | Vulnerable | Species or species habitat may occur within area |
| Phoebetria fusca Sooty Albatross [1075] | Vulnerable | Species or species habitat likely to occur within area |
| Sternula albifrons | | |
| Little Tern [82849] | | Species or species habitat may occur within area |
| <u>Thalassarche bulleri</u> Buller's Albatross, Pacific Albatross [64460] | Vulnerable | Species or species habitat may occur within area |
| <u>Thalassarche cauta</u> Shy Albatross [89224] | Vulnerable* | Foraging, feeding or related behaviour likely to occur within area |
| Thalassarche chrysostoma Grey-headed Albatross [66491] | Endangered | Species or species habitat may occur within area |
| <u>Thalassarche impavida</u> Campbell Albatross, Campbell Black-browed Albatross [64459] | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| Thalassarche melanophris Black-browed Albatross [66472] | Vulnerable | Species or species habitat may occur within |

| Name | Threatened | Type of Presence |
|--|----------------|---|
| | | area |
| Thalassarche salvini | | |
| Salvin's Albatross [64463] | Vulnerable | Foraging, feeding or related behaviour likely to occur |
| Thalassarche steadi | | within area |
| White-capped Albatross [64462] | Vulnerable* | Foraging, feeding or related |
| | | behaviour likely to occur within area |
| Migratory Marine Species | | |
| Balaena glacialis australis | En deu neue d* | On a sing an ana sing habitat |
| Southern Right Whale [75529] | Endangered* | Species or species habitat known to occur within area |
| Caperea marginata | | |
| Pygmy Right Whale [39] | | Species or species habitat may occur within area |
| Carcharodon carcharias | | |
| White Shark, Great White Shark [64470] | Vulnerable | Species or species habitat known to occur within area |
| Caretta caretta | | |
| Loggerhead Turtle [1763] | Endangered | Species or species habitat known to occur within area |
| <u>Chelonia mydas</u> | | |
| Green Turtle [1765] | Vulnerable | Foraging, feeding or related behaviour known to occur within area |
| Dermochelys coriacea | | |
| Leatherback Turtle, Leathery Turtle, Luth [1768] | Endangered | Species or species habitat known to occur within area |
| Lagenorhynchus obscurus | | |
| Dusky Dolphin [43] | | Species or species habitat may occur within area |
| Lamna nasus | | |
| Porbeagle, Mackerel Shark [83288] | | Species or species habitat likely to occur within area |
| Megaptera novaeangliae | | |
| Humpback Whale [38] | Vulnerable | Species or species habitat |
| | | may occur within area |
| Migratory Terrestrial Species | | |
| Hirundapus caudacutus | | |
| White-throated Needletail [682] | Vulnerable | Species or species habitat likely to occur within area |
| Monarcha melanopsis | | |
| Black-faced Monarch [609] | | Species or species habitat likely to occur within area |
| Motacilla flava | | |
| Yellow Wagtail [644] | | Species or species habitat likely to occur within area |
| Myiagra cyanoleuca | | |
| Satin Flycatcher [612] | | Species or species habitat known to occur within area |
| Rhipidura rufifrons | | |
| Rufous Fantail [592] | | Species or species habitat known to occur within area |
| Migratory Wetlands Species | | |
| Actitis hypoleucos | | |
| Common Sandpiper [59309] | | Species or species habitat known to occur within area |

| Name | Threatened | Type of Presence |
|--|-----------------------|--|
| Arenaria interpres | | 0 |
| Ruddy Turnstone [872] | | Species or species habitat known to occur within area |
| Calidris acuminata | | 0 |
| Sharp-tailed Sandpiper [874] | | Species or species habitat known to occur within area |
| Calidris canutus | | . |
| Red Knot, Knot [855] | Endangered | Species or species habitat known to occur within area |
| Calidris ferruginea | | |
| Curlew Sandpiper [856] | Critically Endangered | Species or species habitat known to occur within area |
| Calidris melanotos | | |
| Pectoral Sandpiper [858] | | Species or species habitat likely to occur within area |
| Calidris ruficollis | | |
| Red-necked Stint [860] | | Species or species habitat known to occur within area |
| Calidris subminuta | | 0 |
| Long-toed Stint [861] | | Species or species habitat known to occur within area |
| Calidris tenuirostris | . | . |
| Great Knot [862] | Critically Endangered | Species or species habitat known to occur within area |
| Charadrius bicinctus | | |
| Double-banded Plover [895] | | Species or species habitat known to occur within area |
| Charadrius leschenaultii | | |
| Greater Sand Plover, Large Sand Plover [877] | Vulnerable | Species or species habitat known to occur within area |
| Charadrius mongolus | | |
| Lesser Sand Plover, Mongolian Plover [879] | Endangered | Species or species habitat known to occur within area |
| Gallinago hardwickii | | |
| Latham's Snipe, Japanese Snipe [863] | | Species or species habitat known to occur within area |
| Limicola falcinellus | | |
| Broad-billed Sandpiper [842] | | Species or species habitat known to occur within area |
| Limosa lapponica | | |
| Bar-tailed Godwit [844] | | Species or species habitat known to occur within area |
| Limosa limosa | | |
| Black-tailed Godwit [845] | | Species or species habitat known to occur within area |
| Numenius madagascariensis | | |
| Eastern Curlew, Far Eastern Curlew [847] | Critically Endangered | Species or species habitat known to occur within area |
| Pandion haliaetus | | |
| Osprey [952] | | Species or species habitat likely to occur within area |
| Phalaropus lobatus | | |
| Red-necked Phalarope [838] | | Species or species habitat known to occur within area |

| Name | Threatened | Type of Presence |
|--|------------|--|
| Philomachus pugnax | | |
| Ruff (Reeve) [850] | | Species or species habitat known to occur within area |
| Pluvialis fulva | | |
| Pacific Golden Plover [25545] | | Species or species habitat known to occur within area |
| Pluvialis squatarola | | |
| Grey Plover [865] | | Species or species habitat known to occur within area |
| Tringa brevipes | | |
| Grey-tailed Tattler [851] | | Species or species habitat known to occur within area |
| Tringa glareola | | |
| Wood Sandpiper [829] | | Species or species habitat known to occur within area |
| <u>Tringa nebularia</u> | | |
| Common Greenshank, Greenshank [832] | | Species or species habitat known to occur within area |
| Tringa stagnatilis | | |
| Marsh Sandpiper, Little Greenshank [833] | | Species or species habitat known to occur within area |
| Xenus cinereus | | |
| Terek Sandpiper [59300] | | Species or species habitat known to occur within area |

Other Matters Protected by the EPBC Act

| Listed Marine Species | | [Resource Information] |
|---|-------------------------------|--|
| * Species is listed under a different scientific na | ame on the EPBC Act - Threate | ened Species list. |
| Name | Threatened | Type of Presence |
| Birds | | |
| Actitis hypoleucos | | |
| Common Sandpiper [59309] | | Species or species habitat known to occur within area |
| <u>Anous stolidus</u> | | |
| Common Noddy [825] | | Species or species habitat likely to occur within area |
| Apus pacificus Fork-tailed Swift [678] | | Species or species habitat |
| | | likely to occur within area |
| Ardea alba | | Spacios or aposios habitat |
| Great Egret, White Egret [59541] | | Species or species habitat known to occur within area |
| Ardea ibis | | |
| Cattle Egret [59542] | | Species or species habitat may occur within area |
| Arenaria interpres | | |
| Ruddy Turnstone [872] | | Species or species habitat known to occur within area |
| Calidris acuminata | | |
| Sharp-tailed Sandpiper [874] | | Species or species habitat known to occur within area |

| Name | Threatened | Type of Presence |
|---|-----------------------|--|
| Calidris canutus | | |
| Red Knot, Knot [855] | Endangered | Species or species habitat known to occur within area |
| Calidris ferruginea | | |
| Curlew Sandpiper [856] | Critically Endangered | Species or species habitat known to occur within area |
| Calidris melanotos | | |
| Pectoral Sandpiper [858] | | Species or species habitat likely to occur within area |
| Calidris ruficollis | | |
| Red-necked Stint [860] | | Species or species habitat known to occur within area |
| Calidris subminuta | | |
| Long-toed Stint [861] | | Species or species habitat known to occur within area |
| Calidris tenuirostris | | |
| Great Knot [862] | Critically Endangered | Species or species habitat known to occur within area |
| Charadrius bicinctus | | |
| Double-banded Plover [895] | | Species or species habitat known to occur within area |
| Charadrius leschenaultii | | |
| Greater Sand Plover, Large Sand Plover [877] | Vulnerable | Species or species habitat known to occur within area |
| Charadrius mongolus | | |
| Lesser Sand Plover, Mongolian Plover [879] | Endangered | Species or species habitat known to occur within area |
| Charadrius ruficapillus | | |
| Red-capped Plover [881] | | Species or species habitat known to occur within area |
| Chrysococcyx osculans | | |
| Black-eared Cuckoo [705] | | Species or species habitat known to occur within area |
| Diomedea antipodensis | | |
| Antipodean Albatross [64458] | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| Diomedea epomophora | | |
| Southern Royal Albatross [89221] | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| Diomedea exulans | | |
| Wandering Albatross [89223] | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| Diomedea sanfordi | Endongorod | Europing fooding or related |
| Northern Royal Albatross [64456] | Endangered | Foraging, feeding or related behaviour likely to occur within area |
| <u>Gallinago hardwickii</u> Latham's Snipe, Japanese Snipe [863] | | Species or species habitat known to occur within area |
| Haliaeetus leucogaster | | |
| White-bellied Sea-Eagle [943] | | Species or species habitat known to occur within area |
| Heteroscelus brevipes | | |
| Grey-tailed Tattler [59311] | | Species or species habitat known to occur within area |

| Name | Threatened | Type of Presence |
|--|-----------------------|--|
| Himantopus himantopus | | |
| Pied Stilt, Black-winged Stilt [870] | | Species or species habitat known to occur within area |
| <u>Hirundapus caudacutus</u> White-throated Needletail [682] | Vulnerable | Species or species habitat likely to occur within area |
| Lathamus discolor Swift Parrot [744] | Critically Endangered | Species or species habitat known to occur within area |
| Limicola falcinellus Broad-billed Sandpiper [842] | | Species or species habitat known to occur within area |
| Limosa lapponica Bar-tailed Godwit [844] | | Species or species habitat known to occur within area |
| <u>Limosa limosa</u> Black-tailed Godwit [845] | | Species or species habitat |
| | | known to occur within area |
| Macronectes giganteus Southern Giant-Petrel, Southern Giant Petrel [1060] | Endangered | Species or species habitat may occur within area |
| Macronectes halli | | |
| Northern Giant Petrel [1061] | Vulnerable | Species or species habitat may occur within area |
| Merops ornatus | | |
| Rainbow Bee-eater [670] | | Species or species habitat may occur within area |
| Monarcha melanopsis | | |
| Black-faced Monarch [609] | | Species or species habitat likely to occur within area |
| <u>Motacilla flava</u> Yellow Wagtail [644] | | Species or species habitat likely to occur within area |
| Myiagra cyanoleuca | | |
| Satin Flycatcher [612] | | Species or species habitat known to occur within area |
| Neophema chrysogaster | | |
| Orange-bellied Parrot [747] Numenius madagascariensis | Critically Endangered | Migration route likely to occur within area |
| Eastern Curlew, Far Eastern Curlew [847] | Critically Endangered | Species or species habitat known to occur within area |
| Pachyptila turtur | | |
| Fairy Prion [1066] | | Species or species habitat known to occur within area |
| Pandion haliaetus | | |
| Osprey [952] | | Species or species habitat likely to occur within area |
| Phalaropus lobatus | | |
| Red-necked Phalarope [838] | | Species or species habitat known to occur within area |
| Philomachus pugnax | | One share an exact the bull of the |
| Ruff (Reeve) [850] | | Species or species habitat known to occur within area |
| Phoebetria fusca | | |
| Sooty Albatross [1075] | Vulnerable | Species or species |

| Name | Threatened | Type of Presence |
|---|-------------|--|
| Pluvialis fulva | | habitat likely to occur within area |
| Pacific Golden Plover [25545] | | Species or species habitat known to occur within area |
| <u>Pluvialis squatarola</u> Grey Plover [865] | | Species or species habitat known to occur within area |
| Puffinus carneipes Flesh-footed Shearwater, Fleshy-footed Shearwater [1043] | | Foraging, feeding or related behaviour likely to occur within area |
| Puffinus griseus Sooty Shearwater [1024] | | Species or species habitat may occur within area |
| Recurvirostra novaehollandiae Red-necked Avocet [871] | | Species or species habitat known to occur within area |
| <u>Rhipidura rufifrons</u> Rufous Fantail [592] | | Species or species habitat known to occur within area |
| Rostratula benghalensis (sensu lato) Painted Snipe [889] | Endangered* | Species or species habitat known to occur within area |
| Sterna albifrons Little Tern [813] | | Species or species habitat may occur within area |
| <u>Stiltia isabella</u> Australian Pratincole [818] | | Species or species habitat known to occur within area |
| <u>Thalassarche bulleri</u> Buller's Albatross, Pacific Albatross [64460] | Vulnerable | Species or species habitat may occur within area |
| <u>Thalassarche cauta</u> Shy Albatross [89224] | Vulnerable* | Foraging, feeding or related behaviour likely to occur within area |
| Thalassarche chrysostoma Grey-headed Albatross [66491] | Endangered | Species or species habitat may occur within area |
| Thalassarche impavida Campbell Albatross, Campbell Black-browed Albatross [64459] | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| Thalassarche melanophris Black-browed Albatross [66472] | Vulnerable | Species or species habitat may occur within area |
| <u>Thalassarche salvini</u> Salvin's Albatross [64463] | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| Thalassarche sp. nov. Pacific Albatross [66511] | Vulnerable* | Species or species habitat may occur within area |
| Thalassarche steadi White-capped Albatross [64462] | Vulnerable* | Foraging, feeding or related behaviour likely to occur within area |
| Thinornis rubricollis rubricollis Hooded Plover (eastern) [66726] | Vulnerable | Species or species habitat likely to occur |

| Name | Threatened | Type of Presence |
|---|------------|---|
| | | within area |
| <u>Tringa glareola</u> | | |
| Wood Sandpiper [829] | | Species or species habitat |
| | | known to occur within area |
| <u>Tringa nebularia</u> | | |
| Common Greenshank, Greenshank [832] | | Species or species habitat |
| , L 1 | | known to occur within area |
| | | |
| <u>Tringa stagnatilis</u> Marsh Sandpiper, Little Greenshank [833] | | Species or species habitat |
| Marsh Sandpiper, Little Greenshank [033] | | known to occur within area |
| | | |
| Xenus cinereus | | |
| Terek Sandpiper [59300] | | Species or species habitat |
| | | known to occur within area |
| Mammals | | |
| Arctocephalus forsteri | | |
| Long-nosed Fur-seal, New Zealand Fur-seal [20] | | Species or species habitat |
| | | may occur within area |
| Arctocephalus pusillus | | |
| Australian Fur-seal, Australo-African Fur-seal [21] | | Species or species habitat |
| | | may occur within area |
| - | | |
| Reptiles | | |
| Caretta caretta Loggerhead Turtle [1763] | Endongorod | Species or species habitat |
| Loggernead Turtle [1703] | Endangered | known to occur within area |
| | | |
| <u>Chelonia mydas</u> | | |
| Green Turtle [1765] | Vulnerable | Foraging, feeding or related |
| | | behaviour known to occur within area |
| Dermochelys coriacea | | within area |
| Leatherback Turtle, Leathery Turtle, Luth [1768] | Endangered | Species or species habitat |
| | | known to occur within area |
| | | |
| Whales and other Cetaceans | | [Resource Information] |
| Name | Status | Type of Presence |
| Mammals | | |
| Caperea marginata | | |
| Pygmy Right Whale [39] | | Species or species habitat may occur within area |
| | | may occur within area |
| Delphinus delphis | | |
| Common Dophin, Short-beaked Common Dolphin [60] | | Species or species habitat |
| | | may occur within area |
| Eubalaena australis | | |
| Southern Right Whale [40] | Endangered | Species or species habitat |
| | | known to occur within area |
| | | |
| Lagenorhynchus obscurus Dusky Dolphin [43] | | Species or species habitat |
| Dusky Dolphin [43] | | may occur within area |
| | | |
| <u>Megaptera novaeangliae</u> | | |
| Humpback Whale [38] | Vulnerable | Species or species habitat |
| | | may occur within area |
| Tursiops aduncus | | |
| Indian Ocean Bottlenose Dolphin, Spotted Bottlenose | | Species or species habitat |
| Dolphin [68418] | | likely to occur within area |
| Tursiops truncatus s. str. | | |
| Bottlenose Dolphin [68417] | | Species or species habitat |
| | | may occur within area |
| | | |

Extra Information

| State and Territory Reserves | [Resource Information] |
|---|--|
| Name | State |
| Limeburners Lagoon (Hovells Creek) F.F.R | VIC |
| Serendip Wetlands Education Facility | VIC |
| Regional Forest Agreements | [Resource Information] |
| Note that all areas with completed RFAs have been inc | luded. |
| Name | State |
| West Victoria RFA | Victoria |
| Invasive Species | [Resource Information] |
| Weeds reported here are the 20 species of national sig that are considered by the States and Territories to pos following feral animals are reported: Goat, Red Fox, Ca Landscape Health Project, National Land and Water Re | e a particularly significant threat to biodiversity. The t, Rabbit, Pig, Water Buffalo and Cane Toad. Maps from |
| Name | Status Type of Presence |
| Birds | |
| Acridotheres tristis | |
| Common Myna, Indian Myna [387] | Species or species habitat likely to occur within area |
| Alauda arvensis | |
| Skylark [656] | Species or species habitat likely to occur within area |
| Anas platyrhynchos | |
| Mallard [974] | Species or species habitat likely to occur within area |
| Carduelis carduelis | |
| European Goldfinch [403] | Species or species habitat likely to occur within area |
| Carduelis chloris | |
| European Greenfinch [404] | Species or species habitat likely to occur within area |
| Columba livia | |
| Rock Pigeon, Rock Dove, Domestic Pigeon [803] | Species or species habitat likely to occur within area |
| Passer domesticus | |
| House Sparrow [405] | Species or species habitat likely to occur within area |
| Passer montanus Eurasian Tree Sparrow [406] | Species or species habitat |
| | likely to occur within area |
| Pycnonotus jocosus | |
| Red-whiskered Bulbul [631] | Species or species |
| | |

Name

Streptopelia chinensis Spotted Turtle-Dove [780]

Sturnus vulgaris Common Starling [389]

Turdus merula Common Blackbird, Eurasian Blackbird [596]

Turdus philomelos Song Thrush [597]

Mammals

Canis lupus familiaris Domestic Dog [82654]

Capra hircus Goat [2]

Felis catus Cat, House Cat, Domestic Cat [19]

Lepus capensis Brown Hare [127]

Mus musculus House Mouse [120]

Oryctolagus cuniculus Rabbit, European Rabbit [128]

Rattus norvegicus Brown Rat, Norway Rat [83]

Rattus rattus Black Rat, Ship Rat [84]

Sus scrofa Pig [6]

Vulpes vulpes Red Fox, Fox [18]

Plants

Asparagus asparagoides Bridal Creeper, Bridal Veil Creeper, Smilax, Florist's Smilax, Smilax Asparagus [22473]

Cenchrus ciliaris Buffel-grass, Black Buffel-grass [20213]

Chrysanthemoides monilifera Bitou Bush, Boneseed [18983]

Status

Type of Presence habitat likely to occur within area

Species or species habitat likely to occur within area

Species or species habitat likely to occur within area

Species or species habitat likely to occur within area

Species or species habitat likely to occur within area

Species or species habitat likely to occur within area

Species or species habitat likely to occur within area

Species or species habitat likely to occur within area

Species or species habitat likely to occur within area

Species or species habitat likely to occur within area

Species or species habitat likely to occur within area

Species or species habitat likely to occur within area

Species or species habitat likely to occur within area

Species or species habitat likely to occur within area

Species or species habitat likely to occur within area

Species or species habitat likely to occur within area

Species or species habitat may occur within area

Species or species habitat may occur within area

| Name | Status | Type of Presence |
|--|---------------------|--|
| Chrysanthemoides monilifera subsp. monilifera Boneseed [16905] | | Species or species habitat likely to occur within area |
| Lycium ferocissimum | | |
| African Boxthorn, Boxthorn [19235] | | Species or species habitat likely to occur within area |
| Nassella neesiana | | |
| Chilean Needle grass [67699] | | Species or species habitat likely to occur within area |
| Nassella trichotoma | | |
| Serrated Tussock, Yass River Tussock, Yass Tu Nassella Tussock (NZ) [18884] | Jssock, | Species or species habitat likely to occur within area |
| Rubus fruticosus aggregate | | |
| Blackberry, European Blackberry [68406] | | Species or species habitat likely to occur within area |
| Salix spp. except S.babylonica, S.x calodendror | a & S.x reichardtii | |
| Willows except Weeping Willow, Pussy Willow a Sterile Pussy Willow [68497] | nd | Species or species habitat likely to occur within area |
| Solanum elaeagnifolium | | |
| Silver Nightshade, Silver-leaved Nightshade, Wi Horse Nettle, Silver-leaf Nightshade, Tomato We White Nightshade, Bull-nettle, Prairie-berry, Satansbos, Silver-leaf Bitter-apple, Silverleaf-ne Trompillo [12323] Ulex europaeus | eed, | Species or species habitat likely to occur within area |
| Gorse, Furze [7693] | | Species or species habitat likely to occur within area |
| Nationally Important Wetlands | | [Resource Information] |
| Name | | State |
| Werribee-Avalon Area | | VIC |

Caveat

The information presented in this report has been provided by a range of data sources as acknowledged at the end of the report.

This report is designed to assist in identifying the locations of places which may be relevant in determining obligations under the Environment Protection and Biodiversity Conservation Act 1999. It holds mapped locations of World and National Heritage properties, Wetlands of International and National Importance, Commonwealth and State/Territory reserves, listed threatened, migratory and marine species and listed threatened ecological communities. Mapping of Commonwealth land is not complete at this stage. Maps have been collated from a range of sources at various resolutions.

Not all species listed under the EPBC Act have been mapped (see below) and therefore a report is a general guide only. Where available data supports mapping, the type of presence that can be determined from the data is indicated in general terms. People using this information in making a referral may need to consider the qualifications below and may need to seek and consider other information sources.

For threatened ecological communities where the distribution is well known, maps are derived from recovery plans, State vegetation maps, remote sensing imagery and other sources. Where threatened ecological community distributions are less well known, existing vegetation maps and point location data are used to produce indicative distribution maps.

Threatened, migratory and marine species distributions have been derived through a variety of methods. Where distributions are well known and if time permits, maps are derived using either thematic spatial data (i.e. vegetation, soils, geology, elevation, aspect, terrain, etc) together with point locations and described habitat; or environmental modelling (MAXENT or BIOCLIM habitat modelling) using point locations and environmental data layers.

Where very little information is available for species or large number of maps are required in a short time-frame, maps are derived either from 0.04 or 0.02 decimal degree cells; by an automated process using polygon capture techniques (static two kilometre grid cells, alpha-hull and convex hull); or captured manually or by using topographic features (national park boundaries, islands, etc). In the early stages of the distribution mapping process (1999-early 2000s) distributions were defined by degree blocks, 100K or 250K map sheets to rapidly create distribution maps. More reliable distribution mapping methods are used to update these distributions as time permits.

Only selected species covered by the following provisions of the EPBC Act have been mapped:

- migratory and

- marine

The following species and ecological communities have not been mapped and do not appear in reports produced from this database:

- threatened species listed as extinct or considered as vagrants
- some species and ecological communities that have only recently been listed
- some terrestrial species that overfly the Commonwealth marine area
- migratory species that are very widespread, vagrant, or only occur in small numbers

The following groups have been mapped, but may not cover the complete distribution of the species:

- non-threatened seabirds which have only been mapped for recorded breeding sites

- seals which have only been mapped for breeding sites near the Australian continent

Such breeding sites may be important for the protection of the Commonwealth Marine environment.

Coordinates

-38.03987 144.37434

Acknowledgements

This database has been compiled from a range of data sources. The department acknowledges the following custodians who have contributed valuable data and advice:

-Office of Environment and Heritage, New South Wales -Department of Environment and Primary Industries, Victoria -Department of Primary Industries, Parks, Water and Environment, Tasmania -Department of Environment, Water and Natural Resources, South Australia -Department of Land and Resource Management, Northern Territory -Department of Environmental and Heritage Protection, Queensland -Department of Parks and Wildlife, Western Australia -Environment and Planning Directorate, ACT -Birdlife Australia -Australian Bird and Bat Banding Scheme -Australian National Wildlife Collection -Natural history museums of Australia -Museum Victoria -Australian Museum -South Australian Museum -Queensland Museum -Online Zoological Collections of Australian Museums -Queensland Herbarium -National Herbarium of NSW -Royal Botanic Gardens and National Herbarium of Victoria -Tasmanian Herbarium -State Herbarium of South Australia -Northern Territory Herbarium -Western Australian Herbarium -Australian National Herbarium, Canberra -University of New England -Ocean Biogeographic Information System -Australian Government, Department of Defence Forestry Corporation, NSW -Geoscience Australia -CSIRO -Australian Tropical Herbarium, Cairns -eBird Australia -Australian Government - Australian Antarctic Data Centre -Museum and Art Gallery of the Northern Territory -Australian Government National Environmental Science Program -Australian Institute of Marine Science -Reef Life Survey Australia -American Museum of Natural History -Queen Victoria Museum and Art Gallery, Inveresk, Tasmania -Tasmanian Museum and Art Gallery, Hobart, Tasmania -Other groups and individuals

The Department is extremely grateful to the many organisations and individuals who provided expert advice and information on numerous draft distributions.

Please feel free to provide feedback via the Contact Us page.

© Commonwealth of Australia Department of the Environment GPO Box 787 Canberra ACT 2601 Australia +61 2 6274 1111

Appendix B. PMST Threatened Species Assessment

| Name | EPBC Status | Listing Advice | Likelihood of Impact |
|--|--------------------------|---|---|
| Birds | | | |
| Anthochaera phrygia Regent Honeyeater | Critically Endangered | Foraging, feeding or related behaviour may occur within area | Low – Habitat (woodland/forest) not present. |
| <i>Botaurus poiciloptilus</i> Australasian Bittern | Endangered | Species or species habitat known to occur within area | Low – Habitat (wetland) not present. |
| <i>Calidris canutus</i> Red Knot | Endangered | species or species habitat known to occur within area | Low – Habitat (coastal mud/sand flats) not present. |
| Calidris ferruginea Curlew Sandpiper | Critically Endangered | species or species habitat known to occur within area | Low – Habitat (coastal mud/sand flats) not present. |
| Calidris tenuirostris Great Knot | Critically Endangered | Species or species habitat known to occur within area | Low – Habitat (coastal mud/sand flats) not present. |
| Charadrius leschenaultia Greater Sand Plover | Vulnerable | Species or species habitat known to occur within area | Low – Habitat (coastal mud/sand flats) not present. |
| Charadrius mongolus Lesser Sand Plover | Endangered | Species or species habitat known to occur within area | Low – Habitat (coastal mud/sand flats) not present. |
| Diomedea antipodensis Antipodean Albatross | Vulnerable | Foraging, feeding or related behaviour likely to occur within area | Low – Habitat (marine) not present. |
| Diomedea epomophora Southern Royal Albatross | Vulnerable | Foraging, feeding or related behaviour likely to occur within area | Low – Habitat (marine) not present. |
| Diomedea exulans Wandering Albatross | Vulnerable | Foraging, feeding or related behaviour likely to occur within area | Low – Habitat (marine) not present. |
| Diomedea sanfordi Northern Royal Albatross | Endangered | Foraging, feeding or related behaviour likely to occur within area | Low – Habitat (marine) not present. |
| <i>Grantiella picta</i> Painted Honeyeater | Vulnerable | Species or species habitat known to occur within area | Low – Habitat (woodland/forest) not present. |
| <i>Hirundapus caudacutus</i> White-throated Needletail | Vulnerable | Species or species habitat likely to occur within area | Low – Migratory visitor to Australian airspace, rarely lands. |
| Lathamus discolor Swift Parrot | Critically Endangered | Species or species habitat known to occur within area | Low – Habitat (woodland/forest) not present. |
| Limosa lapponica bauera Bar-tailed Godwit (baueri) | Vulnerable | Species or species habitat known to occur within area | Low – Habitat (coastal mud/sand flats) not present. |
| Limosa lapponica menzbieri Northern Siberian Bar-tailed Godwit | Critically Endangered | Species or species habitat may occur within area | Low – Habitat (coastal mud/sand flats) not present. |
| <i>Macronectes giganteus</i> Southern Giant-Petrel | Endangered | Species or species habitat may occur within area | Low – Habitat (marine) not present. |



| Name | EPBC Status | Listing Advice | Likelihood of Impact |
|--|--------------------------|--|---|
| <i>Macronectes halli</i> Northern Giant Petrel | Vulnerable | Species or species habitat may occur within area | Low – Habitat (marine) not present. |
| Neophema chrysogaster Orange-bellied Parrot | Critically Endangered | Migration route likely to occur within area | Low – Habitat (coastal salt marsh) not present. |
| Numenius madagascariensis Eastern Curlew | Critically Endangered | Species or species habitat known to occur within area | Low – Habitat (coastal mud/sand flats) not present. |
| Pachyptila turtur subantarctica | Vulnerable | Species or species habitat known to occur within area | Low – Habitat (coastal mud/sand flats) not present. |
| Fairy Prion (southern) | | | |
| Pedionomus torquatus Plains-wanderer | Critically Endangered | Species or species habitat likely to occur within area | Low – considered locally extinct. |
| Phoebetria fusca Sooty Albatross | Vulnerable | Species or species habitat likely to occur within area | Low – Habitat (marine) not present. |
| Pterodroma leucoptera leucoptera | Endangered | Species or species habitat may occur within area | Low – Habitat (marine) not present. |
| Gould's Petrel | | | • |
| Rostratula australis Australian Painted Snipe | Endangered | Species or species habitat known to occur within area | Low – Habitat (coastal mud/sand flats) not present. |
| S <i>ternula nereis nereis</i> Australian Fairy Tern | Vulnerable | Species or species habitat known to occur within area | Low – Habitat (coastal mud/sand flats) not present. |
| Thalassarche bulleri Buller's Albatross | Vulnerable | Species or species habitat may occur within area | Low – Habitat (marine) not present. |
| Thalassarche bulleri platei Northern Buller's Albatross | Vulnerable | Species or species habitat may occur within area | Low – Habitat (marine) not present. |
| Thalassarche cauta cauta Shy Albatross | Vulnerable | Foraging, feeding or related behaviour likely to occur within area | Low – Habitat (marine) not present. |
| Thalassarche cauta steadi White-capped Albatross | Vulnerable | Foraging, feeding or related behaviour likely to occur within area | Low – Habitat (marine) not present. |
| Thalassarche chrysostoma Grey-headed Albatross | Endangered | Species or species habitat may occur within area | Low – Habitat (marine) not present. |
| Thalassarche impavida Campbell Albatross | Vulnerable | Foraging, feeding or related behaviour likely to occur within area | Low – Habitat (marine) not present. |
| Thalassarche melanophris Black-browed Albatross | Vulnerable | Species or species habitat may occur within area | Low – Habitat (marine) not present. |
| Thalassarche salvini Salvin's Albatross | Vulnerable | Foraging, feeding or related behaviour likely to occur within area | Low – Habitat (marine) not present. |
| Thinornis rubricollis rubricollis | Vulnerable | Species or species habitat likely to occur within area | Low – Habitat (coastal mud/sand flats) not present. |

| Name | EPBC Status | Listing Advice | Likelihood of Impact |
|--|--------------|---|---|
| Hooded Plover (eastern) | | | |
| Fish | | | |
| Galaxiella pusilla | Vulnerable | Species or species habitat likely to | Low – Habitat |
| Eastern Dwarf Galaxias | Vullerable | occur within area | (aquatic/waterways) not present |
| Prototroctes maraena | Vulnerable | Species or species habitat likely to | Low – Habitat (suitable |
| Australian Grayling | Vullerable | occur within area | aquatic/waterways) not present. |
| Frogs | | | |
| Litoria raniformis | Vulnerable | Species or species habitat known to occur within area | Moderate – May utilise local drains and damp areas for |
| Growling Grass Frog | vullerable | occur within area | dispersal/foraging. |
| nsects | | | |
| Synemon plana | Critically | Species or species habitat known to | Moderate – May utilise disturbed |
| Golden Sun Moth | Endangered | occur within area | tussock grasslands (native or exotic). |
| | | | |
| Mammals | | Species or species habitat known to | |
| Eubalaena australis | Endangered | occur within area | Low – Habitat (marine) not present. |
| Southern Right Whale | | Species or species habitat may occur | law liabitat (marina) nat |
| <i>Megaptera novaeangliae</i> Humpback Whale | Vulnerable | within area | Low – Habitat (marine) not present. |
| | | Foraging, feeding or related | Low – Specie may occasionally |
| Pteropus poliocephalus | Vulnerable | behaviour likely to occur within area | overfly the site, but no feeding o |
| Grey-headed Flying-fox | | | rousting habitat present. |
| Plants | | | |
| Amphibromus fluitans | Vulnerable | Species or species habitat likely to occur within area | Low – Species considered out of range. |
| River Swamp Wallaby-grass | | C | |
| Dianella amoena | Endangered | Species or species habitat may occur within area | Moderate – May occur in disturbed tussock grasslands |
| Matted Flax-lily | | | (native or exotic). |
| Diuris basaltica | Endangered | Species or species habitat may occur within area | Low – Species associated with |
| Small Golden Moths Orchid | | within died | native grasslands. |
| Dodonaea procumbens | Vulnerable | Species or species habitat may occur within area | Low – Species associated with |
| Trailing Hop-bush | | within alea | native grasslands. |
| Glycine latrobeana | Vulnerable | Species or species habitat likely to occur within area | Low – Species associated with |
| Clover Glycine | | | native grasslands. |
| Lachnagrostis adamsonii | | Species or species habitat may occur within area | Moderate – Species may be retained in local drainage |
| Adamson's Blown-grass | Endangered | | channels/roadside drains/wet |
| State of the state | | | areas. |
| Leucochrysum albicans var. tricolor | Endangered | Species or species habitat likely to occur within area | Low – Species associated with |
| Hoary Sunray | Linualiyereu | | native grasslands. |

| Name | EPBC Status | Listing Advice | Likelihood of Impact |
|---|--------------------------|---|---|
| Pimelea spinescens subsp. spinescens | Critically Endangered | Species or species habitat known to occur within area | Moderate – May occur in disturbed tussock grasslands (native or exotic). |
| Plains Rice-flower | | | |
| Prasophyllum frenchii Maroon Leek-orchid | Endangered | Species or species habitat likely to occur within area | Low – Species associated with native grasslands. |
| Pterostylis cucullata Leafy Greenhood | Vulnerable | Species or species habitat may occur within area | Low – Species associated with native grasslands. |
| <i>Rutidosis leptorrhynchoides</i> Button Wrinklewort | Endangered | Species or species habitat likely to occur within area | Low – Species associated with native grasslands. |
| Senecio macrocarpus Large-fruit Fireweed | Vulnerable | Species or species habitat likely to occur within area | Moderate – Species may persist in disturbed grasslands. |
| <i>Xerochrysum palustre</i> Swamp Everlasting | Vulnerable | Species or species habitat may occur within area | Low – Outside of current range. |
| Reptiles | | | |
| <i>Caretta caretta</i> Loggerhead Turtle | Endangered | Species or species habitat known to occur within area | Low – Habitat (marine) not present. |
| <i>Chelonia mydas</i> Green Turtle | Vulnerable | Foraging, feeding or related behaviour known to occur within area | Low – Habitat (marine) not present. |
| <i>Delma impar</i> Striped Legless Lizard | Vulnerable | Species or species habitat known to occur within area | Moderate – May occur in disturbed tussock grasslands (native or exotic). |
| <i>Dermochelys coriacea</i> Leatherback Turtle | Endangered | Species or species habitat known to occur within area | Low – Habitat (marine) not present. |
| <i>Tympanocryptis pinguicolla</i> Grassland Earless Dragon | Endangered | Species or species habitat may occur within area | Low – Thought to be locally extinct. |
| Sharks | | | |
| <i>Carcharodon Carcharias</i> Great White Shark | Vulnerable | Species or species habitat known to occur | Low – Habitat (marine) not present. |